

Affine Histories in Quantum Gravity: Introduction and the Representation for a Cosmological Model

Smaragda Kessari¹

May 2006

The dreams of this work are dedicated to the memory of Nena, Fouskas and Chestnut.

Abstract

It is shown how consistent histories quantum cosmology can be realised through Isham's Histories Projection Operator consistent histories scheme. This is done by using an affine algebra instead of a canonical one and also by using cocycle representations. A regularisation scheme allows us to find a history Hamiltonian which exists as a proper self-adjoint operator. The role of a cocycle choice is also discussed.

1 Introduction to the article

The problem of time in quantum gravity is a particularly complicated and difficult one to understand and deal with. This is mostly because of the fact that general relativity being invariant under the group of diffeomorphisms of the spacetime manifold suggests a type of quantum physics that lacks any notion of time, or at least any notion of time as we know it so far (for reviews see [1] and [2]). Various schemes have been suggested to overcome this problem, one of which involves uncovering an 'internal' time from the existing variables in the theory. In the present article we shall work with this approach in a history context. More precisely, we will discuss the history version of a Friedmann-Robertson-Walker (FRW) universe coupled to a scalar field ϕ , and in which we select (before quantisation) the internal time to be this field ϕ .

Our main concerns are to apply consistent histories theory to quantum gravity and to find the correct representation of the history algebra for quantum cosmology. We will argue that the correct representation for quantum cosmology (with a history Hamiltonian that is self-adjoint) involves using the history analogue of the 'affine group' that has been advocated as the correct canonical group in normal canonical quantum gravity (for an extensive presentation see [14]).

The main reason why the consistent histories approach to quantum theory has been chosen by the author to tackle problems in quantum gravity is the fact that in the consistent

¹s.kessari@ntlworld.com

histories scheme the conventional view of time need not play any fundamental role and as a result—especially in the History Projection Operator (‘HPO’) formalism (see later)—the scheme allows for logical temporal connectives to be defined without prejudice about the nature of time. In this way certain ‘conceptions’ of time can potentially be used in situations where there is no conventional time.

This means that consistent histories might be used naturally and profitably in a theory of quantum gravity where the notion of time as an external parameter disappears. This is particularly true in the case of quantum cosmology where one tries to apply quantum theory to the entire universe. In addition, the conventional quantum-theory split of observer-system—with the associated notions of an ‘external measurement’ and ‘state vector reduction’—is no longer appropriate in quantum cosmology. One of the main features of the consistent history scheme is that ideas of this type play no fundamental role. So that is another good reason for using history ideas in quantum cosmology. We shall mention here that an attempt to apply HPO histories to quantum cosmology has already been made in [13], but in a very different way from that of this paper; in particular, these authors did not use the ‘affine’ history algebra that we shall adopt.

In summary, we will show how the above choices lead to a well-defined Hamiltonian operator in quantum cosmology. We find the ‘history version’ of a quantum cosmological model Hamiltonian in which one uses a history analogue of the affine commutation relations. Then we find a representation of the affine algebra in which the, appropriately regularised, history Hamiltonian exists as a proper renormalised self-adjoint operator.

Of course, this treatment challenges us to apply similar ideas to the notion of space, something that has unfortunately not been researched yet. However, the consistent histories formalism has already generalised the Hilbert space used in quantum theory, even when there was no original intention for that to happen. And one would not be wrong to say that the consistent histories approach allows for the removal of many established prejudices about what is time, space, measurement, observer etc. and lets the theory work without them.

The structure of the article is as follows. In chapter 2 we introduce the reader to the constituent theories and concepts that will later be used to formulate the affine histories theory and the application of the HPO scheme to quantum cosmology. First the basics of consistent histories theory is introduced as it first appeared in the version of Gell-Mann and Hartle [7]. Then the HPO theory is presented, with emphasis on the way in which it can be developed to handle a continuous time parameter [9], [10], [11]. In section 2.2 we introduce the affine group of canonical quantum gravity [15], [16]. In section 2.3 we show how the original canonical quantisation of the FRW cosmological model unintentionally leads to an ‘affine representation’ although that was not appreciated by the authors at that time.

In chapter 3 the novel ‘affine histories’ approach is unravelled. First we just give the basic relations in quantum gravity and then we show how that algebra is realised in quantum cosmology. By doing this we can relate back to the earlier results in [4] for the FRW model plus scalar field. Then we introduce the history analogue of the affine relations and study their representations using coboundaries in the exponential Hilbert space of a simple, but very singular, representation. Finally we introduce a Klauder-type regularisation of the singular products in the definition of the history Hamiltonian. That choice along with a choice of a cocycle lead to our conclusions about the validity of the affine history Hamiltonian.

2 Introduction to the main concepts

2.1 Consistent histories theory

2.1.1 The emergence of histories in standard quantum theory

One way to introduce the histories theory is through the conditional probability scheme of standard quantum theory which is associated with the ideas of state-vector reduction. Suppose an open, Hamiltonian quantum system is subjected to measurements by an external (classical) observer. Let $\hat{U}(t_1, t_0)$ denote the unitary time-evolution operator from time t_0 to time t_1 . If we let $\hat{\rho}(t_0)$ denote the density operator of the quantum system at time t_0 , then at time t_1 , in the Schrödinger representation, it will have evolved through a unitary transformation to

$$\hat{\rho}(t_1) = \hat{U}(t_1, t_0)\hat{\rho}(t_0)\hat{U}(t_1, t_0)^\dagger = \hat{U}(t_1, t_0)\hat{\rho}(t_0)\hat{U}(t_1, t_0)^{-1}. \quad (1)$$

If at time t_1 we make a measurement of a property α_1 , represented by a projection operator $\hat{\alpha}_1$, then the probability that the property will be found is

$$\text{Prob}(\alpha_1 = 1; \hat{\rho}(t_1)) = \text{tr}(\hat{\alpha}_1 \hat{\rho}(t_1)) = \text{tr}(\hat{\alpha}_1(t_1) \hat{\rho}(t_0)), \quad (2)$$

where $\hat{\alpha}_1(t_1)$ is the Heisenberg-picture operator defined as

$$\hat{\alpha}_1(t_1) := \hat{U}(t_1, t_0)^\dagger \hat{\alpha}_1(t_0) \hat{U}(t_1, t_0). \quad (3)$$

Then according to the Von-Neumann-Lüders ‘reduction’ postulate, the density matrix after the measurement is (look, for example, in [3] for more details)

$$\hat{\rho}_{\text{red}}(t_1) := \frac{\hat{\alpha}_1(t_1) \hat{\rho}(t_0) \hat{\alpha}_1(t_1)}{\text{tr}(\hat{\alpha}_1(t_1) \hat{\rho}(t_0))}. \quad (4)$$

If at time $t_2 > t_1$ we perform another measurement on the same system of a second property α_2 (represented by the projection operator $\hat{\alpha}_2$), then the conditional probability of getting $\alpha_2 = 1$ at t_2 given that $\alpha_1 = 1$ at t_1 is

$$\text{Prob}(\alpha_2 = 1, t_2 | \alpha_1 = 1, t_1; \hat{\rho}(t_0)) = \text{tr}(\hat{\alpha}_2(t_2) \hat{\rho}_{\text{red}}(t_1)) = \frac{\text{tr}(\hat{\alpha}_2(t_2) \hat{\alpha}_1(t_1) \hat{\rho}(t_0) \hat{\alpha}_1(t_1))}{\text{tr}(\hat{\alpha}_1(t_1) \hat{\rho}(t_0))}. \quad (5)$$

Then the probability of getting $\alpha_1 = 1$ at t_1 and α_2 at t_2 is (2) $\times$ (5), that is

$$\text{Prob}(\alpha_2 = 1, t_2 \text{ and } \alpha_1 = 1, t_1; \hat{\rho}(t_0)) = \text{tr}(\hat{\alpha}_2(t_2) \hat{\alpha}_1(t_1) \hat{\rho}(t_0) \hat{\alpha}_1(t_1)). \quad (6)$$

Similarly, by performing an n -sequence of measurements, the probability of finding $\alpha_1, \alpha_2, \dots, \alpha_n$ at times $t_1, t_2, \dots, t_n$ is

$$\begin{aligned} \text{Prob}(\alpha_1 = 1, t_1 \text{ and } \alpha_2 = 1, t_2 \text{ and } \dots \alpha_n, t_n; \hat{\rho}(t_0)) \\ = \text{tr}(\hat{\alpha}_n(t_n) \cdots \hat{\alpha}_1(t_1) \hat{\rho}(t_0) \hat{\alpha}_1(t_1) \cdots \hat{\alpha}_n(t_n)). \end{aligned} \quad (7)$$

In the consistent histories approach to quantum theory we call any time-ordered sequence $\alpha := (\hat{\alpha}_1, \hat{\alpha}_2, \dots, \hat{\alpha}_n)$ of projection operators a *homogeneous history*; the associated

set of time points $(t_1, t_2, \dots, t_n)$, with $t_1 < t_2 < \dots < t_n$, is known as the *temporal support* of the history. The main mathematical object of interest in this scheme is the *decoherence functional* $d_\rho(\alpha, \beta)$, which is defined on pairs of homogeneous histories α, β by

$$d_\rho(\alpha, \beta) := \text{tr}(\hat{\alpha}_n(t_n) \cdots \hat{\alpha}_2(t_2) \hat{\alpha}_1(t_1) \hat{\rho}(t_0) \hat{\beta}_1(t_1) \hat{\beta}_2(t_2) \cdots \hat{\beta}_n(t_n)) = \text{tr}(C_{\hat{\alpha}} \rho(t_0) C_{\hat{\beta}}^\dagger), \quad (8)$$

with

$$C_\alpha := \hat{\alpha}_n(t_n) \cdots \hat{\alpha}_2(t_2) \hat{\alpha}_1(t_1) = U(t_0, t_n) \hat{\alpha}_n U(t_n, t_{n-1}) \cdots \hat{\alpha}_2 U(t_2, t_1) \hat{\alpha}_1 U(t_1, t_0) \quad (9)$$

where $\hat{\alpha}_i(t_i) := U(t_i, t_0)^\dagger \hat{\alpha}_i U(t_i, t_0)$ is the Heisenberg picture operator defined with respect to the fiducial time t_0 . The decoherence functional is defined for a closed system, with no external observers and state-vector reductions; it measures the interference between two histories that have some properties at particular times and no reference to measurements is needed. The crucial assumption of the history formalism is that, under certain circumstances, the probability (7) *is* meaningful even in the absence of explicit measurements. The way this is realised is by requiring that $d_\rho(\alpha, \beta) = 0$ for all pairs α, β , $\alpha \neq \beta$, in a set of histories. Such a set of histories is said to be *consistent*.

In this article we will only be interested in the appropriate representation of the histories α and (an analogue of) the ‘class operators’ C_α , leaving the very important subject of the decoherence functional for a later publication. However, it is important to mention some features of the decoherence functional to aid the comprehension of the histories theory and, in particular, its incorporation in quantum cosmology.

One of these is that the decoherence functional encodes all the information about the history—both its dynamical structure and its initial state—whereas the history itself is just an ordered (time-ordered) sequence of projection operators, each one of which is a ‘proposition’ (in the language of quantum logic) that in standard quantum theory refers to the results of measurements at times $t_1, t_2, \dots, t_n$ where $t_1 < t_2 < \dots < t_n$. Let us also clarify here that a proposition asserts the value of some observable in some range of the real numbers at a given time. In our example a history $\alpha = (\hat{\alpha}_1, \hat{\alpha}_2, \dots, \hat{\alpha}_n)$ is the sequential conjunction “ α_1 is true at time t_1 , and then α_2 is true at time t_2 , and then $\dots$, and then α_n is true at time t_n ”.

2.1.2 The Gell-Mann and Hartle approach

In order to understand the approach to history theory that is followed in this article it is useful to describe how it arose originally. The consistent histories approach to quantum theory was discovered independently by Omnès, Griffiths, and Gell-Mann and Hartle (GH) [6], [7]. It has become an area of particular interest in the foundations of quantum theory, and it has provided a new way of both realising and interpreting quantum gravity with its emphasis on the fact that everything that can be said about the physical world (including its classical aspects) can be expressed in terms of sets of decohering histories.

Two main schools in the field of decohering histories are the path-integral approach, followed mainly recently by Hartle and Halliwell [8], and the HPO approach: a generalisation of the GH approach that was discovered and developed by Isham, Linden and Schreckenberg [9], [10]. Here we will only mention the developments that will be relevant to the present article.

Gell-Mann and Hartle made two important suggestions. The first one was that the histories formalism can be extended to include disjoint² *sums* of homogeneous histories, known as *inhomogeneous* histories; this was done by introducing a mechanism for forming a logical ‘or’ for disjoint histories as well as a logical ‘not’.

The second suggestion of GH was that a history could be regarded as a fundamental quantity in its own right and not be just thought of as a time-ordered sequence of projection operators/single-time propositions. They achieved this by introducing a set of axioms that were later reformulated and extended by Isham in order to broaden and give a more solid mathematical foundation to the theory. We shall discuss this shortly.

2.1.3 Histories Projection Operator (HPO) initial conception

The work of Gell-Mann and Hartle gave the history theory a new perspective with new potentials for future research. However, to realise these fully, a new mathematical framework was necessary, and this was provided by Isham [9], and Isham and Linden [10] in the form of the ‘HPO theory’, the consistent histories theory that is used here.

The key step was to introduce ideas of sequential/temporal quantum logic and to represent such propositions by replacing the class operator of GH—which is not a projection operator—with a *tensor* product of projectors—which *is* a projection operator. Note that, until the invention of the HPO formalism, almost all studies of quantum logic had involved propositions at a single time, and were therefore not applicable to the histories theory.

As far as the use of tensor products is concerned, let us first recall that in the GH approach, a (homogeneous) history α is a set $(\hat{\alpha}_1, \hat{\alpha}_2, \dots, \hat{\alpha}_n)$ of single-time propositions which is represented by the class operator $C_\alpha := \hat{\alpha}_n \cdots \hat{\alpha}_2 \hat{\alpha}_1$. The product of projectors is in general not itself a projector (this would require them all to commute pairwise), and hence by using C_α the link with quantum logic is lost. However, the key observation of Isham [9] was that a homogeneous history can also be represented by the tensor product $\hat{D}_\alpha := \hat{\alpha}_n \otimes \hat{\alpha}_{n-1} \otimes \cdots \otimes \hat{\alpha}_1$ which is defined on the *tensor* product of n copies of the single-time Hilbert space $\mathcal{H}$. Unlike C_α , the new quantity $\hat{D}_\alpha$ *is* a projector.

This development fitted in very well with the suggestion by Gell-Mann and Hartle that history theory could be suitable for quantum gravity where the classical notions of space and time are not applicable. As a result, as long as propositions about the analogue of ‘histories’ exist in a theory of this type, it becomes reasonable to look for their representation with projectors on a new ‘history’ Hilbert space.

The HPO scheme has some attractive features. For example, by applying the usual logical operations on projection operators, the space of history propositions can be identified with an orthoalgebra, or lattice³, of history propositions. That way, the set of propositions about the histories becomes equipped with the structure of a non-distributive lattice, and the lattice of projection operators in a suitable Hilbert space can model this generalised-history space.

²Two homogeneous histories $\alpha := (\hat{\alpha}_1, \hat{\alpha}_2, \dots, \hat{\alpha}_n)$ and $\beta := (\hat{\beta}_1, \hat{\beta}_2, \dots, \hat{\beta}_n)$ are *disjoint* if for at least one time point t_i , $\hat{\beta}_i$ is disjoint from $\hat{\alpha}_i$, i.e. the ranges of these two projection operators are orthogonal subspaces of $\mathcal{H}$.

³The notable difference between those two is that in the former the ‘or’ proposition is defined only on pairs of disjoint elements whereas in a lattice it is defined on any pair (therefore, an orthoalgebra seems to be the minimal structure that can be used in quantum logic).

More precisely, the proposition ‘ α_1 at time t_1 ’ can be represented on the two-time history Hilbert space by the operator $\hat{1} \otimes \hat{\alpha}_1$, and similarly the proposition ‘ α_2 at time t_2 ’ can be represented by the operator $\hat{\alpha}_2 \otimes \hat{1}$. Then the sequential conjunction ‘ α_1 at time t_1 and then α_2 at time t_2 ’ is represented by $\hat{1} \otimes \hat{\alpha}_1 \wedge \hat{\alpha}_2 \otimes \hat{1}$, which is equal to $\hat{\alpha}_2 \otimes \hat{\alpha}_1$, the HPO representation of the history $(\hat{\alpha}_1, \hat{\alpha}_2)$. Therefore, a homogeneous history can be identified with a sequential conjunction, which is in agreement with its intrinsic and logical meaning.

2.1.4 Continuous time and the history group in generalised quantum theory and quantum field theory

As our interest is centred in quantum cosmology, the question now arises about how the history formalism can be extended from a finite temporal support to a continuous one and so we need to consider what meaning can be given to ‘continuous’ temporal logic. There have been two approaches to this question by Isham and Linden [10], and by Isham, Linden et al [11]. Both of them have the same starting point, which is the *history group*: a history analogue of the canonical group and the associated canonical commutation relations (CCR) used in single-time quantum theory. This will play a central role in our discussion of quantum cosmology.

The Lie algebra of the Weyl-Heisenberg group of transformations of classical state-space gives the classical observables that can be represented by self-adjoint operators in the associated quantum theory, with the classical Poisson brackets going to operator commutators. For example, consider the quantum theory of a particle moving on the real line $\mathbf{R}$. The Lie algebra of the Weyl-Heisenberg group is⁴

$$[\hat{x}, \hat{p}] = i \quad (10)$$

and what one looks for is an irreducible, unitary representation of this group. Then, according to the Stone and von Neumann theorem, the familiar representation on wave functions $\psi(x)$ is unique up to unitary isomorphisms.

What is sought for in the HPO analogue is a unitary representation of a *history group* with a *history algebra* whose self-adjoint representations will give the projection operators that will represent the propositions about the continuous-time histories of the system.

As discussed in the previous section, a homogeneous history $\alpha = (\hat{\alpha}_1, \hat{\alpha}_2, \dots, \hat{\alpha}_n)$ is represented by the projection operator $\hat{\alpha}_n \otimes \hat{\alpha}_{n-1} \otimes \dots \otimes \hat{\alpha}_1$ that acts on the tensor product $\mathcal{V}_n := \mathcal{H}_n \otimes \mathcal{H}_{n-1} \otimes \dots \otimes \mathcal{H}_1$ of n -copies of the Hilbert space $\mathcal{H}$ of the canonical theory. Since each $\mathcal{H}$ has a representation of the Lie algebra (10), the space $\mathcal{V}_n = \mathcal{H}_n \otimes \mathcal{H}_{n-1} \otimes \dots \otimes \mathcal{H}_1$ carries a representation of the Lie algebra

$$[\hat{x}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}, \hat{x}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}] = [\hat{x}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}, \hat{1} \otimes \hat{x}_{n-1} \otimes \hat{1} \otimes \dots \otimes \hat{1}] = \dots = 0 \quad (11)$$

$$[\hat{p}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}, \hat{p}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}] = [\hat{p}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}, \hat{1} \otimes \hat{p}_{n-1} \otimes \dots \otimes \hat{1}] = \dots = 0 \quad (12)$$

$$[\hat{x}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}, \hat{p}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}] = [\hat{1} \otimes \hat{x}_{n-1} \otimes \dots \otimes \hat{1}, \hat{1} \otimes \hat{p}_{n-1} \otimes \dots \otimes \hat{1}] = \dots = i \quad (13)$$

$$[\hat{x}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}, \hat{1} \otimes \hat{p}_{n-1} \otimes \dots \otimes \hat{1}] = [\hat{1} \otimes \hat{x}_{n-1} \otimes \dots \otimes \hat{1}, \hat{p}_n \otimes \hat{1} \otimes \dots \otimes \hat{1}] = \dots = 0 \quad (14)$$

⁴We set $\hbar=1$ from now on.

etc. More succinctly, we have

$$[\hat{x}_k, \hat{p}_m] = i\delta_{km} \quad (15)$$

$$[\hat{p}_k, \hat{p}_m] = 0 \quad (16)$$

$$[\hat{x}_k, \hat{x}_m] = 0 \quad (17)$$

with $k, m = 1, 2, \dots, n$, where all the operators $\hat{x}_k$ and $\hat{p}_m$ are viewed as acting on the one Hilbert space $\mathcal{V}_n$ of the tensor product of n -copies of the Hilbert space $\mathcal{H} \simeq L^2(\mathbf{R})$ of the canonical theory.

However, what is used in practice, especially in continuous-time histories, is the reverse argument; *i.e.*, one starts from (15)–(17) as defining the history algebra, and then seeks a representation of it. The Stone-Von Neumann theorem shows that there is an essentially unique representation space on $\mathcal{V}_n$. Therefore, each history corresponds to a proposition about the values of position and momentum (or linear combinations of them) at the appropriate times in the temporal support, and is constructed using the spectral projectors of the representation of the history Lie algebra.

In order to construct the continuous-time histories, we suppose that the analogue of (15)–(17) is the algebra

$$[\hat{x}_{t_1}, \hat{p}_{t_2}] = ic_{[t]}\delta(t_1 - t_2) \quad (18)$$

$$[\hat{p}_{t_1}, \hat{p}_{t_2}] = 0 \quad (19)$$

$$[\hat{x}_{t_1}, \hat{x}_{t_2}] = 0 \quad (20)$$

where $-\infty \leq t_1, t_2 \leq \infty$ and $c_{[t]}$ is a dimensional constant which has been introduced to make dimensions on both sides of the equations agree [11]. We note that this algebra is infinite dimensional, as can be seen from the fact that these equations resemble those found in one-dimensional quantum field theory.

Introducing a test-function space, which we assume is some (dense) subset of the space $L^2(\mathbf{R})$ of real, square-integrable functions on $\mathbf{R}$, we can introduce the smeared quantities $\hat{x}_f = \int dt f(t)\hat{x}_t$ and $\hat{p}_g = \int dt g(t)\hat{p}_t$ in terms of which the continuous history algebra (CHA) becomes [11]

$$[\hat{x}_f, \hat{p}_g] = ic_{[t]} \int f(t)g(t) dt = ic_{[t]}(f, g) \quad (21)$$

$$[\hat{x}_{f_1}, \hat{x}_{f_2}] = 0 \quad (22)$$

$$[\hat{p}_{g_1}, \hat{p}_{g_2}] = 0. \quad (23)$$

where $(f, g) := \int f(t)g(t) dt$.

There are two different approaches one can take in order to find the appropriate representation of the history algebra (18)–(20). In the first approach, [10], one tries directly to construct the continuous analogue of a tensor product. This involves using coherent states [10] and leads to the familiar Bosonic Fock space in the guise of an exponential Hilbert space (see below). One can then show that the history group algebra (18)–(20), or the smeared form (21), has a natural representation on this space.

The second approach, [11] starts with the history group, with history algebra (18)–(20), and studies the representations of this infinite-dimensional algebra. The Stone von Neumann theorem does not work in infinite dimensions, and in fact there are infinitely many

unitarily inequivalent representations. Guided by the results in an old QFT paper of Araki [5], one seeks a representation in which the operator that represents history propositions about time-averaged values of energy genuinely exists (Araki showed that, in normal quantum field theory, the requirement that the Hamiltonian for a free field exists, uniquely determines the representation).

We will not discuss any details here of the original papers on continuous history theory as it turns out that the history analogue of the affine algebra has a representation theory that is radically different from that associated with the canonical history algebra in (18)–(20). However, as we shall see, it is still useful to employ the language of Fock spaces and exponential Hilbert spaces, although these entities now arise in a very different way.

However, the fact that the basic history affine algebra representations are very different from those of (18)–(20) is not the only reason why the representation methods in [11] cannot be applied to the affine case. Another major reason is that, in [11], a key role is played by the fact that the Hamiltonian operator along with the configuration variables $\hat{x}_t$ and $\hat{p}_t$ form a (closed) Lie algebra. For if one finds the unitary representations of the associated extended Lie group, then one guarantees the existence of a self-adjoint Hamiltonian. Indeed, this is the scheme used to *construct* the Hamiltonian.

However, as we shall see, in our quantum cosmology model if one adds the Hamiltonian to the set of the affine variables then the resulting collection no longer forms a Lie algebra (this is already true at the Poisson algebra level). As a result, in the affine case one has to try to construct the Hamiltonian directly (from the ‘outset’) as a function of the affine history variables. This is a tricky process involving regularisation and is the procedure which we will follow⁵.

Finally, in order to conceptualise and understand the application of HPO histories theory in cosmology, let us say a few parenthetical words about the application of the CHA in quantum field theory by Isham, Linden et al [11]. We foliate a four-dimensional Minkowski space-time with the aid of a timelike vector n^μ , normalised by $\eta_{\mu\nu}n^\mu n^\nu = 1$ where $\eta_{\mu\nu} = (+, -, -, -)$. The equal time CCR for bosonic quantum field theory are

$$[\phi(\underline{x}_1), \phi(\underline{x}_2)] = 0 = [\pi(\underline{x}_1), \pi(\underline{x}_2)] \quad (24)$$

$$[\phi(\underline{x}_1), \pi(\underline{x}_2)] = i\delta^3(\underline{x}_1 - \underline{x}_2). \quad (25)$$

Following the same steps as we did when going from (10) to (18)–(20), we ‘historise’ the quantum field CCR above and we get

$$[\phi_{t_1}(\underline{x}_1), \phi_{t_2}(\underline{x}_2)] = 0 = [\pi_{t_1}(\underline{x}_1), \pi_{t_2}(\underline{x}_2)] \quad (26)$$

$$[\phi_{t_1}(\underline{x}_1), \pi_{t_2}(\underline{x}_2)] = ic_{[t]}\delta(t_1 - t_2)\delta^3(\underline{x}_1 - \underline{x}_2). \quad (27)$$

One can also rewrite (26)–(27) in a ‘four-vector’ mode as

$$[\phi(X), \phi(Y)] = 0 = [\pi(X), \pi(Y)] \quad (28)$$

$$[\phi(X), \pi(Y)] = ic_{[t]}\delta^4(X - Y) \quad (29)$$

⁵Actually, this phenomenon arises already in normal quantum theory since, apart from the case of the simple harmonic oscillator, the Hamiltonian $\frac{p^2}{2m} + V(x)$ does not combine with x and p to give a Lie algebra. In the field theory analogue, it is only the Hamiltonian of the free field that combines with the other variables to give a Lie algebra.

which however are *not* covariant commutation relations, as the four-vector x_n with $(t, x) \in \mathbf{R} \times \mathbf{R}^3$ (with $n \cdot x_n = 0$) of the covariant theory is related to the space-time point X with $X = tn + x_n$.

2.2 The affine algebra in quantum theory and quantum gravity

2.2.1 The motivation for using affine commutation relations

In [14], and later in [15] and [16], it was extensively and explicitly shown that although the familiar CCR are appropriate for a system whose classical configuration space, Q , is a vector space, this canonical algebra is not appropriate for systems where Q has a different topology. However, group theory ideas *can* still be employed effectively provided that the configuration space is a homogeneous space: *i.e.*, Q has some transitive group, G , of transformations, so that $Q \simeq G/H$ for some subgroup H of G . The canonical group then becomes the semi-direct product $W \times_s G$ where W is a vector space carrying a linear representation of G and such that Q is embedded equivariantly as a G orbit [14]. This new canonical group $W \times_s G$ is often referred to as an ‘affine’ group.

The use of such group-theoretic methods becomes particularly important when the configuration space Q is infinite-dimensional. Experience with standard QFT suggests that the state vectors will be functionals on a ‘distributional dual’ of Q . However, it is not trivial to define what is meant by such a dual when Q is not a vector space (for example, what is meant by a ‘distributional metric’ in quantum gravity?) and the group-theoretical approach to quantisation provides a powerful way of addressing this question.

One simple, but important, example where one has to deal with a non-vector space in quantum theory is the following. Suppose we want to quantise a particle moving on the positive real line. The standard CCR for $Q \simeq \mathbf{R}$ are $[\hat{x}, \hat{p}] = i$ which when exponentiated becomes $e^{-ia\hat{p}}\hat{x}e^{ia\hat{p}} = \hat{x} - a$. But since a is any real number, these CCR are incompatible with the condition $x > 0$. On the other hand, if we instead take the *affine* algebra (the ‘ACR’)

$$[\hat{x}, \hat{\pi}] = i\hat{x}, \quad (30)$$

(where, classically, $\pi = xp$), and if we exponentiate this algebra, then we find that for any (faithful) irreducible representation the spectrum of the operator $\hat{x}$ is either the positive real numbers or the negative real numbers. Indeed, as was first shown by Klauder [17] and then by Pilati [19], there exist two unitarily inequivalent, faithful, irreducible representations of the ACR: one where the spectrum of the variable $\hat{x}$ of the configuration space is strictly greater than zero, and one where it is strictly less than zero. So, by choosing the appropriate representation we are able to quantise the system in a way that is compatible with the classical requirement that $x > 0$.

Similarly, in canonical quantum gravity one wants the metric g_{ij} to be a positive determinant tensor with signature $(+++)$. Therefore, the configuration space, and consequently the unconstrained phase space of (g_{ij}, π^{ij}) , is not a vector space, and the corresponding analogue of the affine algebra (30) is

$$[\hat{g}_{ij}(x), \hat{\pi}_l^k(y)] = \frac{i}{2}(\hat{g}_{il}(x)\delta_j^k + \hat{g}_{jl}(x)\delta_i^k)\delta^{(3)}(x - y) \quad (31)$$

$$[\hat{\pi}_i^j(x), \hat{\pi}_k^l(y)] = \frac{i}{2}(\hat{\pi}_k^j(x)\delta_i^l - \hat{\pi}_i^l(x)\delta_k^j)\delta^{(3)}(x - y) \quad (32)$$

$$[\hat{g}_{ij}(x), \hat{g}_{kl}(y)] = 0. \quad (33)$$

In the simple minisuperspace model of quantum cosmology that is considered below, the variable $g_{ij}(x)$ becomes just the radius R of the universe, and one has to have an algebra that is compatible with the structure of this reduced classical configuration space. But this is nothing but the positive real numbers and hence, as remarked above in the context of (30), the appropriate algebra is

$$[\hat{R}, \hat{\Pi}] = i\hat{R}. \quad (34)$$

2.2.2 The representation of the ACR

As we will be concerned with a history, minisuperspace analogue of (31)–(33) let us consider briefly what is known about the representations of the latter algebra, as discussed in [16].

The group associated with this algebra is the semi-direct product $\mathcal{C} = C^\infty(\Sigma, S(3, \mathbf{R})) \times_s C^\infty(\Sigma, GL^+(3, \mathbf{R}))$ where $S(3, \mathbf{R})$ is the (vector) space of 3×3 real symmetric matrices and Σ is the spatial manifold. When used in canonical quantum gravity, the task is to find unitary representations of this group. Such a unitary representation automatically gives rise to the self-adjoint operators that represent the associated physical variables. From a technical perspective, we can take advantage of the fact that unitary operators are always bounded, unlike the (smeared) operators $\hat{g}(v)$ and $\hat{\pi}(M)$.

One way of finding a representation of the semi-direct product group $\mathcal{C}$ is through the theory of induced representations and Mackey's theorems [21]. However, in such a case *all* the representations found in this way are exhaustive only in the case where $\mathcal{C}$ is a finite-dimensional Lie group. Nevertheless, in [16] some limited results were obtained by treating the infinite-dimensional case as a generalisation of the finite-dimensional one. We shall now briefly discuss some of these results.

If W is a function space $C^\infty(\Sigma, V)$, with V a finite-dimensional vector space (in our example we have $V \simeq S(3, \mathbf{R})$), its (topological) dual space, $C^\infty(\Sigma, V)'$, will play an important role in the quantum theory. Indeed, general spectral theory tells us that, in general, the state vector can always be written as a (possibly vector-space valued) functional $\Psi[\chi]$ of $\chi \in C^\infty(\Sigma, V)'$.

This space, $C^\infty(\Sigma, V)'$, is some set of distributions, and if $\chi \in C^\infty(\Sigma, V)'$ we follow convention and write, rather heuristically, for all $v \in C^\infty(\Sigma, V)$,

$$\chi(v) = \int_\Sigma \chi_{ij}(x) v^{ij}(x) d\theta(x) \quad (35)$$

where $d\theta$ is the measure on the three-manifold Σ associated with some background volume element⁶.

In our case, the simplest example of an element of the distributional dual is the Dirac δ -function object, $\delta_{(x,g)}$, which is defined for each $x \in \Sigma$ and $g \in GL^+(3, (R))$, as

$$\delta_{(x,g)}(v) := g_{ij} v^{ij}(x) \quad (36)$$

for all $v \in C^\infty(\Sigma, S(3, \mathbf{R}))$. Thus we can write

$$\delta_{(x,g)ij}(\cdot) := g_{ij} \delta_x(\cdot) \quad (37)$$

⁶The use of a background volume element can be avoided if one employs densities of an appropriate weight as the test functions v .

where $(\cdot)$ reminds us that this is a generalised function on Σ .

One of the major problems to be dealt with is the existence and properties of the measure μ that needs to be defined on the quantum state domain space $C^\infty(\Sigma, S(3, \mathbf{R}))'$. This is used to give the inner product on the quantum Hilbert space, and it must be chosen so that the representation of the group $\mathcal{C}$ is unitary. However, the space $C^\infty(\Sigma, S(3, \mathbf{R}))'$ is infinite-dimensional, and finding measures of this type is very difficult. On the other hand, the space of distributions of the type in (36) can be identified with the space $\Sigma \times GL^+(3, \mathbf{R})$ via the injection

$$\begin{aligned} \Sigma \times GL^+(3, \mathbf{R}) &\rightarrow C^{\infty'}(\Sigma, GL^+(3, \mathbf{R})) \\ (x, g) &\mapsto g_{ij}\delta_x(\cdot) \end{aligned} \quad (38)$$

On this (rather thin) subspace of distributions we can use the measure $d\theta \otimes dH$ on $\Sigma \times GL^+(3, \mathbf{R})$ where dH is the Haar measure on $GL^+(3, \mathbf{R})$. This enables a representation of $\mathcal{C} = C^\infty(\Sigma, S(3, \mathbf{R})) \times_s C^\infty(\Sigma, GL^+(3, \mathbf{R}))$ to be defined on $L^2(\Sigma \times GL^+(3, \mathbf{R}); d\theta \otimes dH)$ which, taking into account the group law⁷ and $\delta_{(x,g)}(v) := g_{ij}v^{ij}(x)$, is found to be

$$(V(v)\psi)(x, g) = \exp(i\Sigma_{i,j=1}^3 v^{ij}(x)g_{ij})\psi(x, g) \quad (39)$$

$$(U(\Lambda)\psi)(x, g) = \psi(x, \Lambda^{-1}(x)g). \quad (40)$$

We note that this representation of $\mathcal{C}$ is reducible since state functions $\psi(x, g)$ whose support in x is some compact subset $K \subset \mathbf{R}$ form an invariant subspace. This is a problem to which shall return later in our discussion of quantum cosmology.

By expanding (39)–(40) one finds the self-adjoint operators

$$(\hat{g}_{ij}(y)\psi)(x, g) = g_{ij}\delta_x(y)\psi(x, g) \quad (41)$$

$$(\hat{\pi}_i{}^j(y)\psi)(x, g) = -i\delta_x(y)g_{ik}\partial\psi/\partial g_{jk}(x, g). \quad (42)$$

This shows clearly the sense in which the metric operator $\hat{g}_{ij}(x)$ (and the associated momenta $\hat{\pi}_i{}^j(x)$) is concentrated on a single space point. That gave the idea of generalising (37) to sums of distributions like

$$\delta_{(x_1, g^{(1)}; x_2, g^{(2)}; \dots; x_N, g^{(N)})}(\cdot) := \Sigma_{n=1}^N g_{ij}^{(n)} \delta_{x_n}(\cdot) \quad (43)$$

and (39)–(40) are accordingly replaced with

$$(V(v)\psi)(x_1, g^{(1)}; \dots; x_N, g^{(N)}) := \exp(i\Sigma_{n=1}^N \Sigma_{i,j=1}^3 v^{ij}(x_n)g_{ij}^{(n)})\psi(x_1, g^{(1)}; \dots; x_N, g^{(N)}) \quad (44)$$

$$(U(\Lambda)\psi)(x_1, g^{(1)}; \dots; x_N, g^{(N)}) := \psi(x_1, \Lambda^{-1}(x_1)g^{(1)}; \dots; x_N, \Lambda^{-1}(x_N)g^{(N)}). \quad (45)$$

This representation too is reducible. However the possibility arises of taking the direct sum (over N) of all such representations and then introducing a cocycle [22] on the associated exponential Hilbert space. This possibility was raised in [16], and employed by Klauder [18] and Pilati [19]. This suggestion was not developed in [16] but it is one that we shall explore in our history theory.

⁷The group law of this group is, for all $x \in \Sigma$, $(\Lambda_1(x), v_1(x))(\Lambda_2(x), v_2(x)) = (v_1(x) + \Lambda_1(x)v_2(x), \Lambda_1(x)\Lambda_2(x))$ where $\Lambda_1, \Lambda_2 \in C^\infty(\Sigma, GL^+(3, \mathbf{R}))$ and $v_1, v_2 \in C^\infty(\Sigma, S(3, \mathbf{R}))$.

The most serious problem with representations of the type in (44)–(45) is that they are very singular, which causes big problems if one has to try to construct a product of its operators, such as is needed (as we shall see in the following subsection) for the Hamiltonian operator in our quantum cosmological model. This is so first because the relevant product is always calculated at the *same* spatial point (spatial point in the example above, but time point in the cosmological model history case we shall see next) and secondly because of the meaninglessness of a squared Dirac delta function. Klauder [17] suggested a regularisation of the product of such representations which involves ‘dividing by’ a delta function $\delta(0)$ and we shall return to this later in the context of quantum cosmology.

2.3 A Friedmann-Robertson-Walker cosmological model

In [4] the quantisation of the homogeneous and isotropic FRW universe coupled to a scalar field was given. The metric is

$$ds^2 = N(t)^2 dt^2 - R(t)^2 S_{ij} dx^i dx^j \quad (46)$$

where $N(t)$ is the lapse function (the normal component of the deformation vector-hypersurface of constant time), $R(t)$ represents the radius of the universe and S_{ij} is the fixed background metric for a three-space for the curvature constant K taking any of the values 1, 0, -1 for a three-sphere, flat, or hyperbolic universe respectively.

In the case of a massive scalar field the matter Lagrangian is

$$\mathcal{L} = \frac{1}{2}(-\det g)^{1/2}(g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - m^2\phi^2) \quad (47)$$

where ϕ is the scalar field and $g_{\mu\nu}$ is the four-metric of (46). As the spatial metric is homogeneous, there are just three coupled variables: $N(t)$, $R(t)$ and $\phi(t)$.

The ADM (Arnowitt, Deser, Misner) Lagrangian [23] is [4]

$$\mathcal{L} = \pi_R \dot{R} + \pi_\phi \dot{\phi} + N \left(\frac{\pi_R^2}{24R} + 6KR - R^3\phi^2 \frac{m^2}{2} - \frac{\pi_\phi^2}{2R^3} \right) \quad (48)$$

in which N is a Lagrange multiplier, so that the quantity in the parenthesis is set to 0 by the equations of motion. It is common in canonical quantisation to choose one of the variables as an ‘internal time’. In [4] various such choices were made, but here we shall adopt the one that is in line with our affine scheme. This is the choice $\phi = t$. Then, after solving the constraint equation

$$\frac{\pi_R^2}{24R} + 6KR - R^3\phi^2 \frac{m^2}{2} - \frac{\pi_\phi^2}{2R^3} = 0 \quad (49)$$

for π_ϕ , and replacing it in the Lagrangian (48), we are led to the squared Hamiltonian

$$H^2 = \frac{1}{12}R^2\pi_R^2 + 12KR^4 - R^6t^2m^2. \quad (50)$$

In trying to quantise this system, account must be taken of the fact that, classically, R is confined to lie in the interval $(0, \infty)$. However, if we try to quantise the system

in the obvious way (with the wave-function vanishing at the end points) using the usual assignments

$$R \rightarrow R, \quad \pi_R \rightarrow -i \frac{d}{dR} \quad (51)$$

satisfying the CCR

$$[\hat{R}, \hat{\pi}_R] = i \quad (52)$$

then a problem arises because the second operator turns out not to be self-adjoint (although it is Hermitian)[4].

In the original paper [4], this problem was tackled in a rather indirect way by first conjecturing a suitably quantized Hamiltonian for the massless ($m = 0$) and flat three-space ($K = 0$) case, *i.e.*, for the case

$$H^2 = \frac{1}{12} R^2 \pi_R^2 \quad (53)$$

and showing that this has an interpretation as a positive self-adjoint Hamiltonian whose positive square root exists. The operator $\widehat{R\pi_R}$ was then defined as a combination of R and π_R that *is* self-adjoint (and positive) and specifically it was defined that the quantisation of $R^2 \pi_R^2$ is $\frac{d}{dR} R^2 \frac{d}{dR}$ and then $\widehat{R\pi_R}$ was defined to be the square root of this operator. However, we know now from the previous discussion that $R\pi_R$ is classically the ‘affine conjugate’ Π of R , so that the classical Hamiltonian can be written as

$$H^2 = \frac{1}{12} \Pi^2 + 12KR^2 - R^6 t^2 m^2 \quad (54)$$

with the ACR being

$$[\hat{R}, \hat{\Pi}] = i\hat{R} \quad (55)$$

as was discussed in Section 2.2.1. This, physically meaningful, way explains the mathematics and the structure behind the original way that was successfully guessed by Blyth and Isham, and it offers the possibility for further development and applications. The Hamiltonian (54) is the one whose history version we are going to deal with below.

3 Affine histories theory

The logic for the steps that have already been followed was (i) start with the Lie algebra that one *does* have *i.e.*, the one generated by the affine variables R and Π ; (ii) construct the corresponding Lie group; and then (iii) look for the unitary representations of this Lie group. In each such representation the affine variables will arise as self-adjoint operators representing the associated Lie algebra, and so the final step is to construct a self-adjoint Hamiltonian in terms of these affine variables.

We are interested in the history analogue of the ACR which, as we will see, will involve variables $R(t)$ and $\Pi(t)$, and we may start to worry that problems of divergences will arise when we come to construct the history analogue of the Hamiltonian (54). The history analogue of Isham’s or Klauder’s representations and regularisation methods for the affine algebra might be of use here. If these representations do not work, then new representations must be sought. However, we have already argued that those representations are unlikely

to be the correct ones for an infinite-dimensional histories theory, and we shall also see that that those regularisation methods can be replaced with some more physical and efficient ones.

Let us briefly see (i) what *has* been done already in terms of the correct representation; (ii) what does not seem appropriate to be followed; and (iii) what needs to be changed in an ‘affine histories’ scheme.

In [11] a representation of the CHA (18)–(20) was chosen for a particle moving on the real line using the familiar ideas from quantum field theory, especially the use of the bosonic Fock space as the history Hilbert space. This employed the result that the (history analogue of the) Hamiltonian operator and the history variables x_t, p_s form a closed algebra. Then, since the time automorphism was unitarily implementable, the corresponding self-adjoint operator existed and could be identified as the time average of the energy in the history theory. However, we have seen now that the CCR, and consequently the associated history analogue, is not the correct algebra for our quantum cosmology theory where $R > 0$.

On the other hand in [16], and as discussed above, some representations of the ACR in quantum gravity were found using induced representation techniques. However, the metric operator and its conjugate were defined on a single spatial point and an effort to use an extension of that representation leads to its further reducibility. However, it was suggested in [16] that the use of cocycles might change the situation and this suggestion, in its history form, will be studied below.

3.1 Affine history algebra in quantum cosmology

Here, we shall not attempt to find a representation of the whole quantum gravity affine history algebra. Rather, we shall study the relevant history algebra for the quantum cosmological model of interest. So, based on the arguments in [11] that were analysed in Section 2.1.4, we claim that the history version of the ACR (31)–(33) is

$$[\hat{g}_{ij}(x, t), \hat{\pi}_l^k(y, t')] = \frac{i}{2} c_{[t]} (\hat{g}_{il}(x, t) \delta_j^k + \hat{g}_{jl}(x, t) \delta_i^k) \delta^{(3)}(x - y) \delta(t - t') \quad (56)$$

$$[\hat{\pi}_i^j(x, t), \hat{\pi}_k^l(y, t')] = \frac{i}{2} c_{[t]} (\hat{\pi}_k^j(x, t) \delta_i^l - \hat{\pi}_i^l(x, t) \delta_k^j) \delta^{(3)}(x - y) \delta(t - t') \quad (57)$$

$$[\hat{g}_{ij}(x, t), \hat{g}_{kl}(y, t')] = 0 \quad (58)$$

which for a spatially homogeneous geometry becomes

$$[\hat{g}_{ij}(t), \hat{\pi}_l^k(t')] = \frac{i}{2} c_{[t]} (\hat{g}_{il}(t) \delta_j^k + \hat{g}_{jl}(t) \delta_i^k) \delta(t - t') \quad (59)$$

$$[\hat{\pi}_i^j(t), \hat{\pi}_k^l(t')] = \frac{i}{2} c_{[t]} (\hat{\pi}_k^j(t) \delta_i^l - \hat{\pi}_i^l(t) \delta_k^j) \delta(t - t') \quad (60)$$

$$[\hat{g}_{ij}(t), \hat{g}_{kl}(t')] = 0. \quad (61)$$

To be mathematically well-defined, these operators must be smeared with test functions, and we smear $\hat{g}_{ij}(t)$ with a density field $f^{ij}(t)$, and $\hat{\pi}_i^j(t)$ (which is an operator density) with an ordinary field $F_j^i(t)$. Then the ACR (59) become

$$[\hat{g}(f_d), \hat{\pi}(F_o)] = \frac{ic_{[t]}}{2} \int dt \hat{g}_{il}(t) f^{ij}(t) F_j^l(t) + \frac{ic_{[t]}}{2} \int dt \hat{g}_{jl}(t) f^{jl}(t) F_i^l(t) \quad (62)$$

where $\hat{g}(f_d) = \int dt \hat{g}_{ij}(t) f^{ij}(t)$, and $\hat{\pi}(F_o) = \int dt \hat{\pi}_l^k(t) F_k^l(t)$; there is also an appropriate analogue for (60) and (61). The group of the ACR algebra (31)–(33) is the semi-direct product $C^\infty(\Sigma, S(3, \mathbf{R})) \times_s C^\infty(\Sigma, GL^+(3, \mathbf{R}))$, and similarly the group of the algebra (59)–(61) is $C^\infty(T, S(3, \mathbf{R})) \times_s C^\infty(T, GL^+(3, \mathbf{R}))$ where T denotes the ‘time-line’ which could be the whole real line $\mathbf{R}$ or it might be some bounded subset of $\mathbf{R}$ according to the physical situation under consideration.

Let us now consider the FRW cosmological model case analysed in Section 2.3. It is easy to see that the history–ACR will take the form

$$[\hat{R}(t), \hat{\Pi}(t')] = i c_{[t]} \hat{R}(t) \delta(t - t') \quad (63)$$

$$[\hat{\Pi}(t), \hat{\Pi}(t')] = 0, \quad [\hat{R}(t), \hat{R}(t')] = 0 \quad (64)$$

which in smeared form becomes

$$[\hat{R}(f_R), \hat{\Pi}(f_\Pi)] = i c_{[t]} \int dt f_R(t) f_\Pi(t) \hat{R}(t) := i c_{[t]} \hat{R}(f_R f_\Pi) \quad (65)$$

$$[\hat{\Pi}(f_\Pi), \hat{\Pi}(f'_\Pi)] = 0, \quad [\hat{R}(f_R), \hat{R}(f'_R)] = 0 \quad (66)$$

where $\hat{R}(f_R) := \int \hat{R}(t) f_R(t) dt$ and $\hat{\Pi}(f_\Pi) = \int \hat{\Pi}(t) f_\Pi(t) dt$, etc. The corresponding history group is $C(T, \mathbf{R}) \times_s C(T, \mathbf{R}_+) \cong C(T, \mathbf{R} \times_s \mathbf{R}_+)$, *i.e.*, the group of all continuous maps from the time-line T to the group $\mathbf{R} \times_s \mathbf{R}_+$.

The history analogue of the Hamiltonian squared (54) is

$$\hat{H}^2(t) = \frac{1}{12} \hat{\Pi}^2(t) + 12K \hat{R}^4(t) - \hat{R}^6(t) t^2 m^2 \quad (67)$$

and our task is to find a representation of the algebra in (63)–(64) in which $\hat{H}^2(t)$, or a suitably smeared version of it, exists as a proper self-adjoint operator.

Of course, the physically relevant quantity is $\hat{H}$ which will be the square root of (67), and one immediately obvious problem is that when $m \neq 0$, the operator in (67) may well not be positive, and hence its square root cannot be taken. A similar problem arose in the original canonical FRW model and there it was interpreted as a sign that the time choice $t := \phi$ can only be defined in a limited region of the reduced classical state space [4].

We shall return to this issue later, but setting it aside for the moment, our first step is to seek a ‘suitable’ unitary representation of the history group $C(T, \mathbf{R} \times_s \mathbf{R}_+)$ with Lie algebra (63)–(64). By analogy with the discussion in Section 2.2.2, we see that the simplest representation is when the Hilbert space $\mathcal{H}$ is the space of square-integrable functions $\psi(t, r)$, ($r > 0$) with measure $dt dr / r$, and with operators acting as

$$(\hat{r}(s)\psi)(t, r) := \delta(s, t) r \psi(t, r) \quad (68)$$

$$(\hat{\pi}(s)\psi)(t, r) := -i \delta(s, t) r \frac{\partial \psi(t, r)}{\partial r} \quad (69)$$

The (better defined) smeared version is

$$(\hat{r}(f)\psi)(t, r) := f(t) r \psi(t, r) \quad (70)$$

$$(\hat{\pi}(g)\psi)(t, r) := -i g(t) r \frac{\partial \psi(t, r)}{\partial r}. \quad (71)$$

However, this basic representation on $\mathcal{H}$ is inadequate for two reasons:

- (i) It corresponds to a history that is only ‘active’ at a single time-point $t \in T$, whereas we want histories at any finite set of time points $t_1, t_2, \dots, t_N$ for all $N = 1, 2, \dots$. For each fixed N we can do this by taking the (symmetrised) N -fold tensor product of $\mathcal{H}$ with itself, and then we could take the direct sum of these to give the exponential Hilbert space $\mathcal{H}$. However, this turns out to be a very reducible representation of the history algebra, whereas we want one that is irreducible. Adding a cocycle contribution to the action on $\exp \mathcal{H}$ is one possible way of achieving this.
- (ii) If we compute the action of the operator product $\hat{r}(s)\hat{r}(u)$ in $\mathcal{H}$ we find that

$$(\hat{r}(s)\hat{r}(u)\psi)(t, r) = \delta(s, u)\delta(s, t)r^2\psi(t, r) \quad (72)$$

and a similar result for the product $\hat{\pi}(s)\hat{\pi}(u)$. However, the history Hamiltonian in which we are interested is (67) i.e.

$$\hat{H}^2(t) = \frac{1}{12}\hat{\Pi}^2(t) + 12K\hat{R}^4(t) - \hat{R}^6(t)t^2m^2 \quad (73)$$

and it is clear that we are going to have problems while trying to define such singular products of operators at the same time point.

In the case of normal quantum field theory, we know that the representation of the CCR has to be chosen precisely in order that the Hamiltonian should exist, and we might conjecture that the same is true here. In HPO theory we have seen that the Lie algebra (be it canonical or affine) arises on an n -fold tensor product $\mathcal{V}_n = \mathcal{H}_n \otimes \mathcal{H}_{n-1} \otimes \dots \mathcal{H}_1$ of n -copies of the Hilbert space $\mathcal{H}$ of the canonical (resp. affine) theory. For that reason, in the original papers on continuous time, an exponential Hilbert space construction was used as this can be interpreted as a continuous tensor product of copies of the Hilbert space $L^2(\mathbf{R})$ where we have $\mathcal{V}_{\text{cts}} = \otimes_t L_t^2(\mathbf{R}) := \exp L^2(\mathbf{R}, dt)$ [11].

It seems natural therefore to use an exponential Hilbert space in continuous-time history theory. In particular, we wonder if we can use the *cocycle-exponential* Hilbert space construction in order to find a representation of the history affine algebra of our quantum cosmology model in which (73) exists as a well-defined operator.

3.2 Exponential group representations and cocycles

For the present work we shall only need a few basic definitions of cocycles and exponential group representations and these are given briefly here.

First, let $\mathcal{H}$ be a complex Hilbert space. Then the exponential Hilbert space $\exp \mathcal{H}$ is defined by $\exp \mathcal{H} := \mathbf{C} \oplus \mathcal{H} \oplus (\mathcal{H} \otimes_{\text{Symm}} \mathcal{H}) \oplus \dots = \mathbf{C} \oplus_{n=1}^{\infty} \text{Symm}^n \mathcal{H}$, where Symm denotes symmetrised tensor product and where $\mathbf{C}$ is the complex numbers. For the vectors $\phi \in \mathcal{H}$, there are defined the ‘exponential vectors’

$$\exp \phi := 1 \oplus \phi \oplus \frac{1}{\sqrt{2}}\phi \otimes \phi \oplus \frac{1}{\sqrt{3}}\phi \otimes \phi \otimes \phi \oplus \dots = \bigoplus_{n=0}^{\infty} \frac{1}{\sqrt{n}}(\otimes \phi)^n \quad (74)$$

A fundamental property of this construction is that the set of all finite linear combinations of exponential vectors is dense in $\exp \mathcal{H}$: i.e., any vector in $\exp \mathcal{H}$ can be written as the

strong limit of a sequence of finite linear combinations of exponential vectors. Another key result is that the inner products of $\exp \mathcal{H}$ and $\mathcal{H}$ are related by

$$\langle \exp \phi, \exp \psi \rangle_{\exp \mathcal{H}} = e^{\langle \phi, \psi \rangle_{\mathcal{H}}}. \quad (75)$$

Now let $g \rightarrow \hat{A}_g$ be a unitary representation of a topological group G on the Hilbert space $\mathcal{H}$. The *cochains* of different degrees are defined by

$$C^0(G, \mathcal{H}) := \mathcal{H} \quad (76)$$

$$C^q(G, \mathcal{H}) := \{\text{continuous functions from } G \times G \times \dots \times G \text{ to } \mathcal{H}\}, \quad q \geq 1 \quad (77)$$

where the direct product $G \times G \times \dots \times G$ is taken q times. Then the coboundary operator is defined on $C^0(G, \mathcal{H})$, $C^1(G, \mathcal{H})$ and $C^2(G, \mathcal{H})$ as

$$v \in C^0(G, \mathcal{H}), \quad (\delta v)(g) := \hat{A}_g v - v \quad (78)$$

$$f \in C^1(G, \mathcal{H}), \quad (\delta f)(g_1, g_2) := f(g_1) + \hat{A}_{g_1} f(g_2) - f(g_1 g_2) \quad (79)$$

$$f \in C^2(G, \mathcal{H}), \quad (\delta f)(g_1, g_2, g_3) := \hat{A}_{g_1} f(g_2, g_3) - f(g_1 g_2, g_3) + f(g_1, g_2 g_3) - f(g_1, g_2). \quad (80)$$

Using these coboundary operators, the spaces of 1-coboundaries and 1-cocycles, and 2-coboundaries and 2-cocycles are defined respectively as:

$$B^1(G, \mathcal{H}) := \{\beta : G \rightarrow \mathcal{H} \mid \beta(g) = \hat{A}_g v - v\} \quad (81)$$

$$Z^1(G, \mathcal{H}) := \{\beta : G \rightarrow \mathcal{H} \mid \beta(g_1 g_2) = \beta(g_1) + \hat{A}_{g_1} \beta(g_2)\} \quad (82)$$

$$B^2(G, \mathcal{H}) := \{f : G \times G \rightarrow \mathcal{H} \mid f(g_1, g_2) = \beta(g_1) + \hat{A}_{g_1} \beta(g_2) - \beta(g_1 g_2), \beta : G \rightarrow \mathcal{H}\} \quad (83)$$

$$Z^2(G, \mathcal{H}) := \{f : G \times G \rightarrow \mathcal{H} \mid f(g_1, g_2) + f(g_1 g_2, g_3) = f(g_1, g_2 g_3) + \hat{A}_{g_1} f(g_2, g_3)\}. \quad (84)$$

One can easily prove that every coboundary is a cocycle. However, it is not always the case that every cocycle is a coboundary. But *if* it is, then our calculations are much simplified. Since that latter case leads to representations that agree with Klauder's results [17] and [18] we shall in the present article investigate this subcase only. Note however, that we retain the option to follow Klauder's example and choose coboundaries that are singular in some way, so that in effect we do get a non-trivial cohomological situation. Also, as we shall see, an addition of a coboundary to a representation has a marked effect on the regularisation of certain operators that we have to do to get a complete theory.

Let G be a topological group with a unitary representation $g \mapsto \hat{A}_g$ of G on $\mathcal{H}$, and let β be a 1-cocycle. In addition let $\lambda : G \rightarrow U(1)$ be a map such that

$$\lambda(g_1 g_2) = \lambda(g_1) \lambda(g_2) e^{-i \Im \langle \beta(g_1), \hat{A}(g_1) \beta(g_2) \rangle} \quad (85)$$

for all $g_1, g_2 \in G$. Then the associated *exponential representation* of G on $\exp \mathcal{H}$ is the family of operators $\hat{U}_g$, $g \in G$, defined by

$$\hat{U}_g \exp v = \lambda(g) e^{-\frac{1}{2} \|\beta(g)\|^2 - \langle \hat{A}(g)v, \beta(g) \rangle^*} \exp(\hat{A}(g)v + \beta(g)). \quad (86)$$

It can be shown that the operators $\hat{U}_g$ are unitary and that $\hat{U}_{g_1} \hat{U}_{g_2} = \hat{U}_{g_1 g_2}$. Thus we do indeed have a unitary representation of G .

Now suppose that $\beta = \delta\phi_0$ for $\phi_0 \in \mathcal{H}$ i.e., $\beta(g) = \hat{A}_g\phi_0 - \phi_0$ (as follows from (78) and (81)). Then the real function $(g_1, g_2) \mapsto \Im\langle\beta(g_1), \hat{A}(g_1)\beta(g_2)\rangle \in C^2(G, \mathbf{R})$ is the coboundary of $g \rightarrow \Im\langle\hat{A}(g)\phi_0, \phi_0\rangle \in C^1(G, \mathbf{R})$, and we can choose $\lambda(g) = e^{-i\Im\langle\hat{A}(g)\phi_0, \phi_0\rangle}$ for all $g \in G$. Then

$$\hat{U}_g \exp v = e^{-i\Im\langle\hat{A}(g)\phi_0, \phi_0\rangle} e^{-\frac{1}{2}\langle\hat{A}(g)\phi_0 - \phi_0, \hat{A}(g)\phi_0 - \phi_0\rangle - \langle\hat{A}(g)v, \hat{A}(g)\phi_0 - \phi_0\rangle^*} \quad (87)$$

i.e.,

$$\hat{U}_g \exp v = e^{\langle\phi_0, (\hat{A}(g)-1)(\phi_0+v)\rangle} \exp(\hat{A}(g)(v + \phi_0) - \phi_0). \quad (88)$$

Also, if we define

$$E_v(g) = \frac{\langle \exp v, \hat{U}_g \exp v \rangle}{\langle \exp v, \exp v \rangle} \quad (89)$$

then we find that

$$E_v(g) = e^{\langle(\phi_0+v), (\hat{A}(g)-1)(\phi_0+v)\rangle}. \quad (90)$$

3.3 Application to the FRW model

It is clear that equations (88) and (90) can be used in our FRW model form with $\mathcal{H}$ chosen to be $L^2(T \times \mathbf{R}, dt dr/r)$, and with the operators defined in (68) and (69). In particular, we define $\hat{A}(f) := e^{i\hat{r}[f]}$, and the associated exponential representation as being denoted $\hat{U}_f := e^{i\hat{R}[f]}$, where f is a smearing function.

The simplest case for our representation would be to calculate (90) for the case $v = 0$. The exponential vector $\exp 0$ is a cyclic vector and, as a consequence any vector in the Hilbert space can be obtained by taking (the limit of) linear combinations of the form $\hat{U}_g \exp 0$, where g ranges over the elements of the affine history group. In practice, a key role is played by the generating functional

$$E(g) = \langle \exp 0, \hat{U}_g \exp 0 \rangle = e^{\langle\phi_0, (\hat{A}(g)-1)\phi_0\rangle} \quad (91)$$

which for the FRW model case becomes

$$E_R(f) = \langle \exp 0, e^{i\hat{R}[f]} \exp 0 \rangle = e^{\langle\phi_0, (e^{i\hat{r}[f]}-1)\phi_0\rangle} \quad (92)$$

where f is a test function. After computing the functional derivative $\delta/\delta f(s)$ of (92) and then taking the limit $f \rightarrow 0$ we find

$$\langle \exp 0, \hat{R}(s) \exp 0 \rangle = \langle \phi_0, \hat{r}(s) \phi_0 \rangle \quad (93)$$

$$= \int_{\mathbf{R}_+} |\phi_0(s, r)|^2 dr \quad (94)$$

where, in computing the second line, we have used (68).

We now take the second derivative of (92) in the limit $f \rightarrow 0$ and take into account the fact that, on the Hilbert space $\mathcal{H}$, we have

$$(\hat{r}(s)\phi_0)(r, t) = \delta(s-t)r\phi_0(r, t). \quad (95)$$

Moreover, it is easy to show that $\hat{r}(s)\hat{r}(t) = \hat{r}(t)\hat{r}(s)$ and therefore that $\hat{R}(s)\hat{R}(t) = \hat{R}(t)\hat{R}(s)$. As a result we find

$$\langle \exp 0, \hat{R}(s)\hat{R}(t) \exp 0 \rangle = \langle \phi_0, \hat{r}(s)\hat{r}(t)\phi_0 \rangle + \langle \phi_0, \hat{r}(s)\phi_0 \rangle \langle \phi_0, \hat{r}(t)\phi_0 \rangle \quad (96)$$

$$= \delta(s-t) \int_{\mathbf{R}_+} |\phi_0(s, r)|^2 r dr + \int_{\mathbf{R}_+} |\phi_0(s, r_1)|^2 dr_1 \int_{\mathbf{R}_+} |\phi_0(t, r_2)|^2 dr_2. \quad (97)$$

Similarly

$$E_{\Pi}(f) := \langle \exp 0, e^{i\hat{\Pi}[f]} \exp 0 \rangle = e^{\langle \phi_0, (e^{i\hat{\pi}[f]} - 1)\phi_0 \rangle} \quad (98)$$

and we find

$$\langle \exp 0, \hat{\Pi}(s) \exp 0 \rangle = \langle \phi_0, \hat{\pi}(s)\phi_0 \rangle \quad (99)$$

$$= -i \int_{\mathbf{R}_+} \phi_0(s, r)^* \frac{\partial \phi_0(s, r)}{\partial r} dr \quad (100)$$

and

$$\langle \exp 0, \hat{\Pi}(s)\hat{\Pi}(t) \exp 0 \rangle = \langle \phi_0, \hat{\pi}(s)\hat{\pi}(t)\phi_0 \rangle + \langle \phi_0, \hat{\pi}(s)\phi_0 \rangle \langle \phi_0, \hat{\pi}(t)\phi_0 \rangle. \quad (101)$$

3.3.1 The action on $\exp 0$

Moving beyond expectation values we note that (88) with $v = 0$ gives

$$\hat{U}_g \exp 0 = e^{\langle \phi_0, (\hat{A}_g - 1)\phi_0 \rangle} \exp\{(\hat{A}_g - 1)\phi_0\} \quad (102)$$

which for the FRW model case becomes

$$e^{i\hat{R}[f]} \exp 0 = e^{\langle \phi_0, (e^{i\hat{r}[f]} - 1)\phi_0 \rangle} \exp\{(e^{i\hat{r}[f]} - 1)\phi_0\}. \quad (103)$$

After we perform exponential expansion and take the limit $\frac{\delta}{\delta f(s)}|_{f \rightarrow 0}$ we find:

$$\hat{R}(s) \exp 0 = \langle \phi_0, \hat{r}(s)\phi_0 \rangle \exp 0 + \hat{r}(s)\phi_0 \quad (104)$$

where the ‘+’ on the right hand side refers to the sum in the exponential Hilbert space: *i.e.*, the direct sum of the symmetrised tensor products.

After taking the second derivative of (103), in the limit $f \rightarrow 0$ we find

$$\begin{aligned} \hat{R}(s)\hat{R}(t) \exp 0 &= \langle \phi_0, \hat{r}(s)\hat{r}(t)\phi_0 \rangle \exp 0 + \langle \phi_0, \hat{r}(s)\phi_0 \rangle \langle \phi_0, \hat{r}(t)\phi_0 \rangle \exp 0 \\ &+ \langle \phi_0, \hat{r}(s)\phi_0 \rangle \hat{r}(t)\phi_0 + \langle \phi_0, \hat{r}(t)\phi_0 \rangle \hat{r}(s)\phi_0 + \hat{r}(s)\hat{r}(t)\phi_0 \\ &+ \frac{1}{\sqrt{2}} \{ \hat{r}(s)\phi_0 \otimes \hat{r}(t)\phi_0 + \hat{r}(t)\phi_0 \otimes \hat{r}(s)\phi_0 \}. \end{aligned} \quad (105)$$

Similarly, we find

$$\begin{aligned} \hat{\Pi}(s)\hat{\Pi}(t) \exp 0 &= \langle \phi_0, \hat{\pi}(s)\hat{\pi}(t)\phi_0 \rangle \exp 0 + \langle \phi_0, \hat{\pi}(s)\phi_0 \rangle \langle \phi_0, \hat{\pi}(t)\phi_0 \rangle \exp 0 \\ &+ \langle \phi_0, \hat{\pi}(s)\phi_0 \rangle \hat{\pi}(t)\phi_0 + \langle \phi_0, \hat{\pi}(t)\phi_0 \rangle \hat{\pi}(s)\phi_0 + \hat{\pi}(s)\hat{\pi}(t)\phi_0 \\ &+ \frac{1}{\sqrt{2}} \{ \hat{\pi}(s)\phi_0 \otimes \hat{\pi}(t)\phi_0 + \hat{\pi}(t)\phi_0 \otimes \hat{\pi}(s)\phi_0 \}. \end{aligned} \quad (106)$$

where

$$(\hat{\pi}(s)\phi_0)(r, t) = -i\delta(s - t)r\frac{\partial\phi_0}{\partial r}(r, t). \quad (107)$$

In a similar way we find

$$\begin{aligned} \hat{R}(s)\hat{\Pi}(t)\exp 0 &= \langle\phi_0, \hat{r}(s)\hat{\pi}(t)\phi_0\rangle + \langle\phi_0, \hat{r}(s)\phi_0\rangle\hat{\pi}(t)\phi_0 \\ &+ \langle\phi_0, \hat{\pi}(t)\phi_0\rangle\hat{r}(s)\phi_0 + \hat{r}(s)\hat{\pi}(t)\phi_0 \end{aligned} \quad (108)$$

and similarly for $\hat{\Pi}(\lambda)\hat{R}(s)$.

It is easy to check that the affine algebra of the operators $\hat{R}(s)$ and $\hat{\Pi}(s)$ does indeed obey the classical Poisson brackets, as it should; *i.e.*,

$$\begin{aligned} [\hat{R}(s), \hat{\Pi}(t)]\exp 0 &= \hat{R}(s)\hat{\Pi}(t)\exp 0 - \hat{\Pi}(t)\hat{R}(s)\exp 0 \\ &= \langle\phi_0, [\hat{r}(s), \hat{\pi}(t)]\phi_0\rangle\exp 0 + [\hat{r}(s), \hat{\pi}(t)]\phi_0 \\ &= i\delta(s - t)\hat{R}(s)\exp 0 \end{aligned} \quad (109)$$

and also by performing a third order expansion of the exponentials it is easy to check that

$$[\hat{R}(s), \hat{\Pi}(t)\hat{\Pi}(m)]\exp 0 = i\delta(s - m)\hat{\Pi}(t)\hat{R}(s)\exp 0 + i\delta(s - t)\hat{R}(s)\hat{\Pi}(m)\exp 0 \quad (110)$$

and that

$$[\hat{\Pi}(s), \hat{R}(t)\hat{R}(m)]\exp 0 = -i\delta(s - m)\hat{R}(t)\hat{R}(m)\exp 0 - i\delta(s - t)\hat{R}(t)\hat{R}(m)\exp 0 \quad (111)$$

and that, indeed, all the classical Poisson brackets between $\hat{R}(s)$ and $\hat{\Pi}(s)$ are respected.

From equations (101), (105), (95), (96), (107) and (106) it is easy to see that the quantities $\hat{R}(s)^2$ and $\hat{\Pi}(s)^2$ can *not* be evaluated within the present Hilbert space (and therefore within the present exponential Hilbert space), just as the square of the Dirac delta function can not be defined, or just as the Dirac delta function with a zero argument does not have any meaning (it diverges). This is true for any choice of the coboundary, and therefore the need for some sort of regularisation becomes apparent.

3.3.2 The $\hat{\tau}$ operator

A key ingredient of the proposed regularisation is the introduction of the family of operators, $\hat{\tau}(s)$, $s \in T$, which are defined by

$$(\hat{\tau}(s)\psi)(t, r) := \delta(s - t)\psi(t, r). \quad (112)$$

In smeared form we have

$$(\hat{\tau}(f)\psi)(t, r) := f(t)\psi(t, r). \quad (113)$$

It is useful also to introduce the operators $\hat{r}$ and $\hat{\pi}$ defined on $L^2(T \times \mathbf{R}_+; dt dr/r)$ as

$$(\hat{r}\psi)(t, r) := r\psi(t, r) \quad (114)$$

$$(\hat{\pi}\psi)(t, r) := -ir\frac{\partial\psi(t, r)}{\partial r} \quad (115)$$

which obey the ACR $[\hat{r}, \hat{\pi}] = i\hat{r}$. One can easily prove that, for all $s, s' \in T$,

$$(\hat{\tau}(s)\hat{\tau}(s')\psi)(t, r) = \delta(s - s')(\hat{\tau}(s)\psi)(t, r), \quad (116)$$

so that $\hat{\tau}(s)\hat{\tau}(s') = \delta(s - s') \hat{\tau}(s)$. We also have the commutation relations

$$[\hat{\tau}(s), \hat{\tau}(s')] = 0 \quad (117)$$

$$[\hat{\tau}(s), \hat{r}] = 0 \quad (118)$$

$$[\hat{\tau}(s), \hat{\pi}] = 0. \quad (119)$$

The critical point is that the new operator $\hat{\tau}(s)$ is related to the operators $\hat{r}(s)$ and $\hat{\pi}(s)$ by

$$\hat{r}(s) = \hat{\tau}(s)\hat{r} \quad (120)$$

$$\hat{\pi}(s) = \hat{\tau}(s)\hat{\pi}. \quad (121)$$

This will be used extensively in what follows.

3.3.3 $\hat{\tau}(\chi)$ as a projection operator

One can prove that the operator $\hat{\tau}(s)$ is a type of projection operator, as can intuitively be seen from (116) or from the relation

$$\hat{\tau}(s)\hat{\tau}(s') = \delta(s - s')\hat{\tau}(s). \quad (122)$$

More precisely, for $\hat{\tau}(f) = \int \hat{\tau}(s)f(s)ds$ and f any test function, we have, using equation (122),

$$\hat{\tau}(f)\hat{\tau}(h) = \hat{\tau}(fh) \quad (123)$$

which shows that (123) actually corresponds to a representation of the *ring* structure of the space of test functions $C(T, \mathbf{R})$. Moreover, if χ is any characteristic function of a subset of T , then from (123) we have

$$\hat{\tau}(\chi)\hat{\tau}(\chi) = \hat{\tau}(\chi^2) = \hat{\tau}(\chi) \quad (124)$$

which means that $\hat{\tau}(\chi)$ is a genuine *projection* operator.

It now makes sense to define the operator $\hat{t}$ by

$$(\hat{t}\psi)(t, r) := t\psi(t, r), \quad (125)$$

which is self-adjoint on $L^2(T \times \mathbf{R}_+; dt dr/r)$. Making use of the spectral theorem we have that, for all t in T and all Borel functions f ,

$$(f(\hat{t})\psi)(t, r) = f(t)\psi(t, r) = (\hat{\tau}(f)\psi)(t, r). \quad (126)$$

In other words,

$$\hat{\tau}(f) = f(\hat{t}) \quad (127)$$

for all t in T .

Note that it is clear from (122) that the unsmeared quantity $\hat{\tau}(s)$ is not itself a projection operator since, formally, we have

$$\hat{\tau}(s)\hat{\tau}(s) = \delta(0)\hat{\tau}(s) \quad (128)$$

and, of course, the right hand side of this expression is not well-defined because of the $\delta(0)$ factor. However, motivated by ideas discussed at length by Klauder, the form of

(128) suggests that it might be useful to define a *regularised* form of the operator product according to the definition⁸

$$\hat{\tau}(t)\hat{\tau}(t)|_{reg} := \hat{\tau}(t) \quad (129)$$

for each t in T . We shall see the utility of this construction shortly.

3.3.4 The operator $\hat{p}_t$

The reducibility of the representation on $\mathcal{H}$ of the history affine algebra can be understood in terms of (118) and (119) which show that for all test functions $\hat{f}$, $\hat{\tau}(f)$ commutes with the operators $\hat{r}(s)$ and $\hat{\pi}(s)$.

To explore this situation further, let us assume for now on that $T = \mathbf{R}$ for our history group $C(T, \mathbf{R} \times_s \mathbf{R}_+)$ with Lie algebra (63)–(64). Then, one can define a conjugate operator $\hat{p}_t = -id/dt$ which satisfies

$$[\hat{t}, \hat{p}_t] = i \quad (130)$$

and which has the associated one-parameter family of unitary operators

$$\hat{v}(s) := e^{is\hat{p}_t} \quad (131)$$

where $s \in \mathbf{R}$. We see at once that, for all $s \in \mathbf{R}$,

$$\hat{v}(s) \hat{t} \hat{v}(s)^{-1} = \hat{t} + s \quad (132)$$

and the representation of the extended algebra on $\mathcal{H} \simeq L^2(\mathbf{R} \times \mathbf{R}_+; dt dr/r)$ is now *irreducible*. The complete set of commutation relations that is associated with this irreducible representation is

$$[\hat{r}(t), \hat{r}(t')] = 0 \quad (133)$$

$$[\hat{r}(t), \hat{\pi}(t')] = i\delta(t - t')\hat{r}(t) \quad (134)$$

$$[\pi(t), \pi(t')] = 0 \quad (135)$$

supplemented with

$$\hat{v}(s)\hat{r}(t)\hat{v}(s)^{-1} = \hat{r}(t + s) \quad (136)$$

$$\hat{v}(s)\hat{\pi}(t)\hat{v}(s)^{-1} = \hat{\pi}(t + s) \quad (137)$$

for all s, t in $\mathbf{R}$.

3.4 The regularisation procedure

3.4.1 The basic definition

At the end of section 3.3.2 we observed that the new operator $\hat{\tau}(s)$, $s \in \mathbf{R}$, is related to the operators $\hat{r}(s)$ and $\hat{\pi}(s)$ by

$$\hat{r}(s) = \hat{\tau}(s)\hat{r} \quad (138)$$

$$\hat{\pi}(s) = \hat{\tau}(s)\hat{\pi}. \quad (139)$$

⁸In order to keep the dimensions matching up, we should really introduce a constant d whose dimension is time, and then define $\hat{\tau}(t)\hat{\tau}(t)|_{reg} := d\hat{\tau}(t)$ rather than (129). However, in what follows we will ignore such niceties. If necessary, the dimensions can always be corrected at the end of the calculations.

These relations, together with $\hat{\tau}(t)\hat{\tau}(s) = \delta(t-s)\hat{\tau}(s)$, give as a consequence that

$$[\hat{r}(t), \hat{\pi}(s)] = [\hat{r}, \hat{\pi}]\hat{\tau}(t)\hat{\tau}(s) = i\hat{r}\delta(t-s)\hat{\tau}(t) = i\delta(t-s)\hat{r}(t) \quad (140)$$

so that, in this particular representation, the ‘ultralocal’ factor $\delta(t-s)$ in the affine-history algebra comes from the product of $\hat{\tau}(t)$ with $\hat{\tau}(s)$.

The definition (129) of the regularised product $\hat{\tau}(t)\hat{\tau}(t)|_{reg} := \hat{\tau}(t)$ then suggests that if f is any real-valued function of $\hat{r}$, a regularised form of the function $f(\hat{r}(t))$ of $\hat{r}(t)$ can be defined by⁹

$$f(\hat{r}(t))_{reg} := f(\hat{r})\hat{\tau}(t) \quad (141)$$

for each $t \in \mathbf{R}$. Similarly,

$$h(\hat{\pi}(t))_{reg} := h(\hat{\pi})\hat{\tau}(t) \quad (142)$$

for any function h of $\hat{p}$. Here, $f(\hat{r})$ and $h(\hat{\pi})$ are defined in the usual way using the spectral theorem for self-adjoint operators. As a result, these regularised products have the property that

$$[f(\hat{r}(t))_{reg}, h(\hat{\pi}(s))_{reg}] = [f(\hat{r}), h(\hat{\pi})]\delta(t-s)\hat{\tau}(t). \quad (143)$$

3.4.2 The Hamiltonian for the case $m = 0$ and $K = 0$; action on $\exp 0$

Let us start with the simplest case, which is when the mass, m , and curvature, K , parameters both vanish. Then the history Hamiltonian (squared) is formally

$$\hat{H}^2(t) = \frac{1}{12}\hat{\Pi}^2(t) \quad (144)$$

and our task is to give some mathematical meaning to this expression, and to its integrals over regions of the time axis.

Note that, as we said earlier, a critical requirement for any regularisation scheme is that the regularised form of the right hand side of (144) must be a *positive* self-adjoint operator in order that the square root can be taken to give the operator which we are actually seeking, namely $\hat{H}(t)$. Of course, we might expect that the taking of this square root may itself involve some sort of regularisation.

We will explore this matter for the exponential Hilbert space quantisation discussed in Section 3.2 and with a coboundary ϕ_0 that is a function of t and r . Whether or not the function ϕ_0 is a proper element of $L^2(\mathbf{R} \times \mathbf{R}_+, dt dr/r)$ —and hence a genuine coboundary—remains to be seen.

To see the type of regularisation scheme that is suggested, let us begin with equation (106) and note that, formally,

$$\begin{aligned} \hat{\Pi}(t)^2 \exp 0 &= \langle \phi_0, \hat{\pi}(t)^2 \phi_0 \rangle \exp 0 + \langle \phi_0, \hat{\pi}(t) \phi_0 \rangle^2 \exp 0 \\ &+ \langle \phi_0, \hat{\pi}(t) \phi_0 \rangle \hat{\pi}(t) \phi_0 + \hat{\pi}(t)^2 \phi_0 + \sqrt{2} \hat{\pi}(t) \phi_0 \otimes \hat{\pi}(t) \phi_0 \end{aligned} \quad (145)$$

The only divergent terms here are $\langle \phi_0, \hat{\pi}(t)^2 \phi_0 \rangle$ and $\hat{\pi}(t)^2 \phi_0$, and we can regularise these using the techniques introduced above to give

$$\langle \phi_0, \hat{\pi}(t)_{reg}^2 \phi_0 \rangle := \langle \phi_0, \hat{\pi}^2 \hat{\tau}(t) \phi_0 \rangle \quad (146)$$

⁹If one wants to be very precise it is necessary to take care of the dimensions with the add of the additional dimensional constant c mentioned earlier.

and

$$\hat{\pi}(t)_{reg}^2 \phi_0 = \hat{\pi}^2 \hat{\tau}(t) \phi_0. \quad (147)$$

This suggests the definition

$$\begin{aligned} \hat{\Pi}(t)^2|_{reg} \exp 0 &:= \langle \phi_0, \hat{\pi}^2 \hat{\tau}(t) \phi_0 \rangle \exp 0 + \langle \phi_0, \hat{\pi}(t) \phi_0 \rangle^2 \exp 0 \\ &+ \langle \phi_0, \hat{\pi}(t) \phi_0 \rangle \hat{\pi}(t) \phi_0 + \hat{\pi}^2 \hat{\tau}(t) \phi_0 + \sqrt{2} \hat{\pi}(t) \phi_0 \otimes \hat{\pi}(t) \phi_0. \end{aligned} \quad (148)$$

In particular, the expectation value of $\hat{H}(t)_{reg}^2$ in the state $\exp 0$ is

$$\langle \hat{H}^2(t)_{reg} \rangle = \frac{1}{12} \left(\langle \phi_0, \hat{\pi}^2 \hat{\tau}(t) \phi_0 \rangle + \langle \phi_0, \hat{\pi}(t) \phi_0 \rangle^2 \right). \quad (149)$$

Now, as remarked above, the positivity of (149) is a necessary condition if $\langle \hat{H}^2(t)_{reg} \rangle$ is to be the expectation value of a positive operator, and hence one whose square root can be taken. However, we note that, since $\hat{\pi}$ is self-adjoint, $\langle \phi_0, \hat{\pi}^2 \hat{\tau}(t) \phi_0 \rangle = \langle \hat{\pi} \hat{\phi}_0, \hat{\pi} \hat{\tau}(t) \phi_0 \rangle = \langle \hat{\pi} \hat{\phi}_0, \hat{\tau}(t) \hat{\pi} \phi_0 \rangle$ where we have also used the fact that $[\hat{\pi}, \hat{\tau}(t)] = 0$. It follows that the right hand side of (149) is non-negative provided that

$$\langle \hat{\pi} \hat{\phi}_0, \hat{\tau}(t) \hat{\pi} \phi_0 \rangle \geq 0. \quad (150)$$

However, we have that $(\hat{\tau}(t) \phi_0)(s, r) = \delta(t - s) \phi_0(s, r)$ and hence

$$\langle \hat{\pi} \hat{\phi}_0, \hat{\pi} \hat{\tau}(t) \phi_0 \rangle = \int_{\mathbf{R}_+} (\hat{\pi} \phi_0)^*(r, t) (\hat{\pi} \phi_0)(r, t) \frac{dr}{r} \quad (151)$$

and the right hand side is indeed greater or equal to zero since $\hat{\pi}$ is a self-adjoint operator on $L^2(\mathbf{R}_+, dr/r)$. Thus $\langle \hat{H}^2(t)_{reg} \rangle$ is greater than or equal to zero for any choice of the coboundary ϕ_0 .

3.4.3 The Hamiltonian for the case $m = 0$ and $K = 0$; action on $\exp v$

The results above involving $\exp 0$ are encouraging, but of course the fact that the single matrix element $\langle \exp 0, \hat{H}^2(t)_{reg} \exp 0 \rangle$ is non-negative is not sufficient to guarantee that the same can be said for the operator $\hat{H}^2(t)_{reg}$ as a whole. To do this, we need to study the matrix elements $\langle \exp u, \hat{H}^2(t)_{reg} \exp v \rangle$ for arbitrary u, v in $\mathcal{H}$; or, to be more precise, we need to *define* the regularised operator $\hat{H}^2(t)_{reg}$ by giving regularised values for these matrix elements.

The calculations are more complicated than those just involving $\exp 0$ but the basic results are easy to understand, and here we will just give the main ones. One key result is that for any $u, v \in \mathcal{H}$, we have

$$\begin{aligned} \hat{\Pi}(s) \hat{\Pi}(t) \exp v &= \langle (\phi_0 + v), \hat{\pi}(s) \hat{\pi}(t) (\phi_0 + v) \rangle \exp v \\ &+ \langle (\phi_0 + v), \hat{\pi}(s) (\phi_0 + v) \rangle \langle (\phi_0 + v), \hat{\pi}(t) (\phi_0 + v) \rangle \exp v \\ &+ \langle (\phi_0 + v), \hat{\pi}(s) (\phi_0 + v) \rangle \hat{\pi}(t) (\phi_0 + v) \\ &+ \langle (\phi_0 + v), \hat{\pi}(t) (\phi_0 + v) \rangle \hat{\pi}(s) (\phi_0 + v) + \hat{\pi}(s) \hat{\pi}(t) (\phi_0 + v) \\ &+ \frac{1}{\sqrt{2}} \{ \hat{\pi}(s) (\phi_0 + v) \otimes \hat{\pi}(t) (\phi_0 + v) + \hat{\pi}(t) (\phi_0 + v) \otimes \hat{\pi}(s) (\phi_0 + v) \} \end{aligned} \quad (152)$$

and also

$$\begin{aligned} \frac{\langle \exp u, \hat{\Pi}(t) \hat{\Pi}(s) \exp v \rangle}{\langle \exp u, \exp v \rangle} &= \langle (\phi_0 + u), \hat{\pi}(t) \hat{\pi}(s) (\phi_0 + v) \rangle \\ &+ \langle (\phi_0 + u), \hat{\pi}(t) (\phi_0 + v) \rangle \langle (\phi_0 + u), \hat{\pi}(s) (\phi_0 + v) \rangle. \end{aligned} \quad (153)$$

We see at once that the regularised matrix elements of $\hat{H}^2(t)$ are (or, to be more precise, can consistently be defined to be)

$$\begin{aligned} \frac{\langle \exp u, \hat{H}^2(t)_{reg} \exp v \rangle}{\langle \exp u, \exp v \rangle} &= \frac{1}{12} \left\{ \langle (\phi_0 + u), \hat{\pi}^2 \hat{\tau}(t) (\phi_0 + v) \rangle \right. \\ &+ \left. \langle (\phi_0 + u), \hat{\pi}(t) (\phi_0 + v) \rangle \langle (\phi_0 + u), \hat{\pi}(t) (\phi_0 + v) \rangle \right\}. \end{aligned} \quad (154)$$

In particular, for any u in $\mathcal{H} \simeq L^2(\mathbf{R} \times \mathbf{R}_+; dt dr/r)$ we have

$$\begin{aligned} \frac{\langle \exp u, \hat{H}^2(t)_{reg} \exp u \rangle}{\langle \exp u, \exp u \rangle} &= \frac{1}{12} \left\{ \langle \hat{\pi}(\phi_0 + u), \hat{\tau}(t) \hat{\pi}(\phi_0 + u) \rangle \right. \\ &+ \left. \langle (\phi_0 + u), \hat{\pi}(t) (\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{\pi}(t) (\phi_0 + u) \rangle \right\}. \end{aligned} \quad (155)$$

The analogue of the argument used above in the context of (151) shows that the right hand side of (154) is non-negative for any choice of coboundary ϕ_0 . Thus, as desired, the regularised operator $\hat{H}^2(t)_{reg}$ is non-negative. Clearly, the choice of ϕ_0 affects the ‘ground state’ energy of the history system.

3.4.4 The definition of $\hat{H}(t)$ for the case $m = 0$ and $K = 0$

We must consider how to construct a regularised operator $\hat{H}_{reg}(t)$ from the given regularised operator $\hat{H}^2(t)_{reg}$. As an example of the basic ideas, consider on the starting space $\mathcal{H}$ the formal operator $\hat{h}^2(t) := \frac{1}{12} \hat{\pi}(t)^2$ which is divergent, but whose regularised form would be defined as

$$\hat{h}^2(t)_{reg} := \frac{1}{12} \hat{\pi}^2 \hat{\tau}(t). \quad (156)$$

One cannot literally take the square root of the operator in (156) since there is no operator $\hat{v}(t)$ such that $\hat{v}(t)^2 = \hat{\tau}(t)$. What is needed is to use the same regularisation technique as before. In particular we note that $\hat{\pi}^2$ is a positive, self-adjoint operator on $L^2(\mathbf{R}_+, dr/r)$ and hence it has a square root $\sqrt{\hat{\pi}^2}$. This suggests defining an operator

$$\hat{h}(t)_{reg} := \frac{1}{\sqrt{12}} \sqrt{\hat{\pi}^2} \hat{\tau}(t) \quad (157)$$

on $\mathcal{H} = L^2(\mathbf{R} \times \mathbf{R}_+, dt dr/r)$. Clearly, we have the relation

$$(\hat{h}(t)_{reg})^2 = \hat{h}^2(t)_{reg}. \quad (158)$$

These ideas can be developed to construct an operator $\hat{H}(t)_{reg}$ on $\exp \mathcal{H}$ starting with the operator $\hat{H}^2(t)_{reg}$. The details of this will be given in a later paper that will deal with the decoherence function in this history quantum cosmology model.

3.4.5 The Hamiltonian for the case $m \neq 0$ and $K \neq 0$; action on $\exp 0$

For general values of the parameters m and K , the calculations are far more complicated. The history Hamiltonian is formally

$$\hat{H}^2(t) = \frac{1}{12}\hat{\Pi}^2(t) + 12K\hat{R}^4(t) - \hat{R}^6(t)t^2m^2 \quad (159)$$

and the obvious problem here is that one sees immediately that this operator is not necessarily positive for $m \neq 0$, and hence its square root cannot be taken.

To investigate this matter let us start with the simplest expectation value to study, which is

$$\langle \exp 0, \hat{H}^2(t) \exp 0 \rangle = \frac{1}{12} \left(\langle \phi_0, \hat{\pi}(t)^2 \phi_0 \rangle + \langle \phi_0, \hat{\pi}(t) \phi_0 \rangle^2 \right) + 12KA(t) - m^2t^2B(t) \quad (160)$$

where $A(t) := \langle \exp 0, \hat{R}^4(t) \exp 0 \rangle$ and $B(t) := \langle \exp 0, \hat{R}^6(t) \exp 0 \rangle$. In computing these latter expressions we use the basic formula (92) which we functionally differentiate the appropriate number of times. As a result we find that the regularised expression of (160) has the form

$$\langle \exp 0, \hat{H}^2(t)_{reg} \exp 0 \rangle = \frac{1}{12} \left(\langle \phi_0, \hat{\pi}^2 \hat{\tau}(t) \phi_0 \rangle + \langle \phi_0, \hat{\pi} \hat{\tau}(t) \phi_0 \rangle^2 \right) + 12KA(t)_{reg} - m^2t^2B(t)_{reg} \quad (161)$$

where

$$\begin{aligned} A(t)_{reg} &= \langle \phi_0, \hat{r}^4 \hat{\tau}(t) \phi_0 \rangle + \langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle^4 + 4\langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle \langle \phi_0, \hat{r}^3 \hat{\tau}(t) \phi_0 \rangle \\ &+ 3\langle \phi_0, \hat{r}^2 \hat{\tau}(t) \phi_0 \rangle^2 + 6\langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle \langle \phi_0, \hat{r}^2 \hat{\tau}(t) \phi_0 \rangle \langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle \end{aligned} \quad (162)$$

and

$$\begin{aligned} B(t)_{reg} &= \langle \phi_0, \hat{r}^6 \hat{\tau}(t) \phi_0 \rangle + \langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle^6 + 10\langle \phi_0, \hat{r}^3 \hat{\tau}(t) \phi_0 \rangle^2 + 15\langle \phi_0, \hat{r}^2 \hat{\tau}(t) \phi_0 \rangle^3 \\ &+ 6\langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle \langle \phi_0, \hat{r}^5 \hat{\tau}(t) \phi_0 \rangle + 15\langle \phi_0, \hat{r}^2 \hat{\tau}(t) \phi_0 \rangle \langle \phi_0, \hat{r}^4 \hat{\tau}(t) \phi_0 \rangle \\ &+ 15\langle \phi_0, \hat{r}^4 \hat{\tau}(t) \phi_0 \rangle \langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle^2 + 15\langle \phi_0, \hat{r}^2 \hat{\tau}(t) \phi_0 \rangle \langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle^4 \\ &+ 60\langle \phi_0, \hat{r}^3 \hat{\tau}(t) \phi_0 \rangle \langle \phi_0, \hat{r}^2 \hat{\tau}(t) \phi_0 \rangle \langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle \\ &+ 16\langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle^2 \langle \phi_0, \hat{r}^3 \hat{\tau}(t) \phi_0 \rangle + 49\langle \phi_0, \hat{r}^2 \hat{\tau}(t) \phi_0 \rangle^2 \langle \phi_0, \hat{r} \hat{\tau}(t) \phi_0 \rangle^2. \end{aligned} \quad (163)$$

After using the definition (112) we arrive at a squared regularised Hamiltonian whose precise form depends strongly on the coboundary $\phi_0(r, t)$. We note that for some choices of the function $\phi_0(r, t)$ the quantities $A(t)$ and $B(t)$ will diverge even though the coboundary is a member of $L^2(\mathbf{R} \times \mathbf{R}_+; dt dr/r)$. As an illustration of a family of cocycles where this does not happen consider $\phi_0(r, t) := Cr^{n_1}e^{-ar^{n_2}-ibr^{n_3}}$, where $\{n_1, n_2, n_3\}$ are integers, $\{a, b\} \in \mathbf{R}_+$ and C is a (real) constant. We confirmed (with the aid of Maple) that for different choices of a, b, n_1, n_2, n_3 and C all the parts of the Hamiltonian have different values, which means that the Hamiltonian is very ‘cocycle dependant’.

This is important for several reasons. One reason is that the negative contribution to the squared Hamiltonian can have different values. This is important since the physical Hamiltonian is the square root of the right hand side of (160) which should, therefore, be a positive operator; however, its positivity depends on the value of the last negative term.

One relevant remark here is that, unlike normal ordering, the τ -regularisation technique we have employed is such as to maintain the heuristic positivity of operators. Thus, for example, it is clear the the right hand side of (163) is positive for any t and all choices of ϕ_0 for which the relevant integrals over r are finite.

If our conclusions about using the ‘internal’ time $t = \phi$ were similar in history theory to the analogous one that arises in the normal canonical scheme, we would say that the formalism can only be used in situations where t is sufficiently small that the matrix elements of $\hat{H}^2(t)_{reg}$ are positive. To discuss this in the present situation let us use the following example.

Examples

We will take some specific examples for several functions of the cocycle to see what the above expectation value of the squared Hamiltonian become in practice; in particular we shall see how the negative term of (161) appears and for which functions ϕ_0 it can take a ‘small enough’ value so as to guarantee the positivity of the quantity (if any) whose square root we wish to take.

First we integrate (161) over t using (112). This gives (for real-valued cocycles)

$$\begin{aligned} \langle \exp 0, \hat{H}^2(t)_{reg} \exp 0 \rangle = & -\frac{1}{12} \int_0^\infty \phi_0(r, t) \left[\frac{\partial}{\partial r} \phi_0(r, t) + r \frac{\partial^2}{\partial r^2} \phi_0(r, t) \right] dr \\ & - \frac{1}{12} \left[\int_0^\infty \phi_0(r, t) \frac{\partial}{\partial r} \phi_0(r, t) dr \right]^2 + 12KA(t) - m^2 t^2 B(t) \end{aligned} \quad (164)$$

where

$$\begin{aligned} A(t) = & \int_0^\infty \phi_0(r, t)^2 r^3 dr + 3 \left[\int_0^\infty \phi_0(r, t)^2 r dr \right]^2 + 4 \int_0^\infty \phi_0(r, t)^2 dr \int_0^\infty \phi_0(r, t)^2 r^2 dr \\ & + 6 \left[\int_0^\infty \phi_0(r, t)^2 dr \right]^2 \int_0^\infty \phi_0(r, t)^2 r dr + \left[\int_0^\infty \phi_0(r, t)^2 dr \right]^4 \end{aligned} \quad (165)$$

and there is an analogous expression for $B(t)$.

As a demonstration of how the choice of a cocycle affects the results, let us take the simple example where $\phi_0(r, t) := Cre^{-r^n}$, where C is a (real) constant, and n is an integer¹⁰. In this case, with the aid of Maple, we arrive at the conclusion that the range of the possible values for t increases with n and with, for example, $t^2 < 34.3/m^2$ (approximately) for $K = 1$, $n = 100$, $C < 10^{-2}$. We get physical results with all values $\{-1, 0, 1\}$, although whether the results for $K = -1$ are physical (i.e., give a range of t values for which the left hand side of (164) is positive) depends on the value of n , unlike the ones for $K = 1$ which are always acceptable.

What makes a big difference in the range of the t values is the value of the constant a if we modify our example to one with $\phi_0(r, t) := Cre^{-ar^n}$. In this case big values of a give a much wider range for t : for example, for $a = 10^6$, $n = 1$, $K = -1, 0, 1$, $C < 10^{-2}$ we find $t^2 < 5.3 \times 10^{32}/m^2$. It is clear from these results that the range of t over which the expected value of the history FRW Hamiltonian makes sense is very cocycle dependant.

¹⁰We have made the simple choice $\phi_0(r, t) = Cre^{-r^n}$ as this lies in the domain of the self-adjoint operator $\hat{\pi}(f)$, and it makes the expectation value $\langle \phi_0, \hat{\pi}(f) \phi_0 \rangle$ vanish, which simplifies the calculations a little. Note that this particular class of functions ϕ_0 is not square-integrable in t . Nevertheless, it gives rise to meaningful expressions for the regularised operator $\hat{H}^2(t)_{reg}$.

3.4.6 The Hamiltonian for the case $m \neq 0$ and $K \neq 0$; action on $\exp u$

For the general case of $u \neq 0$ where u is any function in $\mathcal{H} \simeq L^2(\mathbf{R} \times \mathbf{R}_+; dt dr/r)$ we can perform calculations analogous to those in the previous section and we arrive at the following expression:

$$\begin{aligned} \frac{\langle \exp u, \hat{H}^2(t)_{reg} \exp u \rangle}{\langle \exp u, \exp u \rangle} &= \frac{1}{12} \left(\langle (\phi_0 + u), \hat{\pi}^2 \hat{\tau}(t)(\phi_0 + u) \rangle + \langle (\phi_0 + u), \hat{\pi} \hat{\tau}(t)(\phi_0 + u) \rangle^2 \right) \\ &+ 12K A_u(t)_{reg} - m^2 t^2 B_u(t)_{reg} \end{aligned} \quad (166)$$

where

$$\begin{aligned} A_u(t)_{reg} &= \langle (\phi_0 + u), \hat{r}^4 \hat{\tau}(t)(\phi_0 + u) \rangle + \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle^4 \\ &+ 4 \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{r}^3 \hat{\tau}(t)(\phi_0 + u) \rangle + 3 \langle (\phi_0 + u), \hat{r}^2 \hat{\tau}(t)(\phi_0 + u) \rangle^2 \\ &+ 6 \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{r}^2 \hat{\tau}(t)(\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle \end{aligned} \quad (167)$$

and

$$\begin{aligned} B_u(t)_{reg} &= \langle (\phi_0 + u), \hat{r}^6 \hat{\tau}(t)(\phi_0 + u) \rangle + \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle^6 \\ &+ 10 \langle (\phi_0 + u), \hat{r}^3 \hat{\tau}(t)(\phi_0 + u) \rangle^2 + 15 \langle (\phi_0 + u), \hat{r}^2 \hat{\tau}(t)(\phi_0 + u) \rangle^3 \\ &+ 6 \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{r}^5 \hat{\tau}(t)(\phi_0 + u) \rangle \\ &+ 15 \langle (\phi_0 + u), \hat{r}^2 \hat{\tau}(t)(\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{r}^4 \hat{\tau}(t)(\phi_0 + u) \rangle \\ &+ 15 \langle (\phi_0 + u), \hat{r}^4 \hat{\tau}(t)(\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle^2 \\ &+ 60 \langle (\phi_0 + u), \hat{r}^3 \hat{\tau}(t)(\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{r}^2 \hat{\tau}(t)(\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle \\ &+ 16 \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle^2 \langle (\phi_0 + u), \hat{r}^3 \hat{\tau}(t)(\phi_0 + u) \rangle \\ &+ 49 \langle (\phi_0 + u), \hat{r}^2 \hat{\tau}(t)(\phi_0 + u) \rangle^2 \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle^2 \\ &+ 15 \langle (\phi_0 + u), \hat{r}^2 \hat{\tau}(t)(\phi_0 + u) \rangle \langle (\phi_0 + u), \hat{r} \hat{\tau}(t)(\phi_0 + u) \rangle^4. \end{aligned} \quad (168)$$

We see that our conclusions will be of essentially the same qualitative type as those in the previous section where $u = 0$.

Our first general conclusion is that, when working with cocycles, the regularisation technique we have employed means that the choice of cocycle has a significant effect on the final results. In general, if we construct two cocycle representations of the history algebra with cocycles that differ by a coboundary, then the two representations are unitarily equivalent. This means that if we take a representation whose cocycle *is* a coboundary, then this representation is unitarily equivalent to the representation in which the cocycle is not present at all (i.e., it is equal to zero) and one might expect therefore that nothing new can be obtained by using only a coboundary. However, as we have seen, the form of the regularised squared-Hamiltonian depends significantly on the coboundary. Thus specifying the coboundary is an important ingredient in determining the final, regularised form of the history Hamiltonian.

4 Conclusions

We have made a preliminary study of the history approach to canonical quantum gravity for the model of an FRW universe coupled to a scalar field ϕ . We have taken an approach in which we choose an internal time before quantisation rather than the other alternatives in which, for example, one looks at a history analogue of the Wheeler-DeWitt equation [12]. The particular choice we have made is $t = \phi$ and we are able to compare our history approach with the much earlier canonical study of this situation [4].

An important part of our programme was to use a history analogue of the affine commutation relations for the metric field rather than the normal canonical quantities. This is important as it helps to guarantee that the quantised metric is indeed a metric. However, this introduces many novel mathematical problems as the representations of the history analogue of the affine commutators are not at all known, whereas for normal canonical fields the history analogues look like conventional free bosonic fields, and the standard Fock space can be employed.

We started with the very singular representation of the history ACR on $L^2(\mathbf{R} \times \mathbf{R}; dt dr / r)$ in which only a single time point is probed. We have extended this to an arbitrary finite number of time points by going to an exponential Hilbert space with a coboundary. Then we have introduced a Klauder-type regularisation method for the (squared) Hamiltonian and shown how the final regularised energy-squared depends on the choice of the coboundary. It turned out that when $m \neq 0$, the problem with positivity of this operator does not arise for us quite as it did in the old canonical quantisation scheme. In particular, we found that there exist (many) choices for the cocycle which can give a positive value for the expected squared Hamiltonian for a wide range of t values; different cocycle-choices give very different results for the expectation value of the Hamiltonian.

Many interesting tasks await future research. Firstly it would be interesting to look at representations with cocycles that are not coboundaries to see how this affects our final results, particularly the positivity of the operators. This involves some very complicated calculations which could be done, for example, by using Maple. Then we have to show how the square roots can be taken for the massless case, and for the appropriate ranges of t for the massive case. Finally, in this line we should look for some more representations of the history affine algebra and try to find a complete theory. This will be important when we try to extend the scheme to more complicated minisuperspace models, or to the full quantum gravity theory.

Then, beyond all this, we must study the decoherence functional and look for interesting consistent sets of histories.

Acknowledgements

I want to thank Chris Isham for many very lengthy and very useful discussions. I also want to thank Charles Wang for introducing me to Maple programming and Peter McClintock for his careful reading through the paper.

I want to thank the future that let this work happen.

References

- [1] C. J. Isham, in ‘Integrable Systems, Quantum Groups, and Quantum Field Theories’, ed. L. A. Ibort and M. A. Rodriguez, London, Kluwer Academic Publishers, 157 (1993); gr-qc/9210011.
- [2] C. J. Isham, in ‘Canonical Quantum Gravity: From Classical to Quantum’, ed. J. Ehlers. H. Friedrich, Springer-Verlag.
- [3] C. J. Isham, ‘Quantum Theory, Mathematical and Structural Foundations’, Imperial College Press (1995).
- [4] W. F. Blyth and C. J. Isham, Phys. Rev. D, **11**(4), 768 (1975).
- [5] H. Araki, J. Math. Phys. **1**, 492 (1960).
- [6] R. B. Griffiths, J. Stat. Phys. **36**, 219 (1984); R. Omnès, Rev. Mod. Phys., **64**, 339 (1992).
- [7] M. Gell Mann and J. B. Hartle, Phys. Rev. D, **47**, 3345 (1993); in ‘Complexity, Entropy and the Physics of Information’, ed. W. H. Zurek, Addison-Wesley, Reading, Vol. VIII, 425 (1990).
- [8] J. Halliwell and M. Ortiz, Phys. Rev. D, **48**, 748 (1993); J. J. Halliwell, Phys. Rev. D, **48**, 2739 (1993); Phys. Rev. D, **48**, 4785 (1993); J. Hartle, Phys. Rev. D, **44**, 3173 (1991); J. Hartle, in Proceedings on the 1992 Les Houches School, Gravitation and Quantisation (1993); M. Gell Mann and J. B. Hartle, Phys. Rev. D, **47**, 3345 (1993).
- [9] C. J. Isham, J. Math. Phys. **35**(5), 2157 (1994).
- [10] C. J. Isham and N. Linden, J. Math. Phys. **35**(10), 5452 (1994); C. J. Isham, N. Linden and S. Schreckenberg, J. Math. Phys. **35**(12), 6360 (1994).
- [11] C. J. Isham and N. Linden, J. Math. Phys. **36**(10), 5392 (1995); C. J. Isham, N. Linden, K. Savvidou and S. Schreckenberg, J. Math. Phys. **39**(4), 1818 (1998).
- [12] J. J. Halliwell and P. Wallden Phys. Rev. D, **73**, 2401 (2006).
- [13] C. Anastopoulos and N. Savvidou, Class. Quantum Grav. **22**, 1841 (2005).
- [14] C. J. Isham, ‘Topological and global aspects of quantum theory’ in ‘Relativity, groups and topology II’, Les Houches summer school, Session XL, 27 Juin–4 Août 1983, édité par Bryce S. DeWitt et Raymond Stora, Elsevier Science Publishers B.V. (1984).
- [15] C. J. Isham and A. C. Kakas, Class. Quantum Grav. **1**, 621 (1984).
- [16] C. J. Isham and A. C. Kakas, Class. Quantum Grav. **1**, 633 (1984).
- [17] J. R. Klauder, Phys. Rev. D, **2**(2), 272 (1970); in ‘Relativity’, ed. M. S. Carmeli, S. I. Flicker and L. Witten (New York: Plenum) (1970); in ‘Lectures in Theoretical Physics’, Volume XIVB, ed. W. E. Brittin, Colorado Associated University Press, Boulder, Colorado, (1973); Annals of Physics, NY **79**, 111 (1973).

- [18] J. Klauder, J. Math. Phys. **11**, 609 (1970).
- [19] M. Pilati, Phys. Rev. D, **26**(10), 2645 (1982); Phys. Rev. D, **28**(4), 729 (1983).
- [20] G. Francisco and M. Pilati, Phys. Rev. D, **31**(2), 241 (1985).
- [21] G. W. Mackey, ‘Unitary Group Representations in Physics, Probability and Number Theory’, New York: Benjamin Cummings (1978).
- [22] J. Erven and B. J. Falkowski, ‘Low order cohomology and Applications, Lecture notes in Mathematics’, **877**, Berlin: Springer (1981).
- [23] R. Arnowitt S. Deser, C. W. Misner, Phys. Rev. **113**, 745 (1959); **116**, 1322 (1959); **117**, 1595 (1960); ‘Gravitation: an introduction to current research’, ed. L. Witten (1962).